arXiv:1405.0713v2 [math.CO] 26 Aug 2015

Further result on acyclic chromatic index of planar graphs

Tao Wang [a, b, *] Yaqiong Zhang [b]

[a]Institute of Applied Mathematics
Henan University, Kaifeng, 475004, P. R. China

[b]College of Mathematics and Information Science
Henan University, Kaifeng, 475004, P. R. China

July 1, 2015

Abstract

An acyclic edge coloring of a graph G is a proper edge coloring such that every cycle is colored with at least three colors. The acyclic chromatic index $\chi'_a(G)$ of a graph G is the least number of colors in an acyclic edge coloring of G. It was conjectured that $\chi'_a(G) \leq \Delta(G) + 2$ for any simple graph G with maximum degree $\Delta(G)$. In this paper, we prove that every planar graph G admits an acyclic edge coloring with $\Delta(G) + 6$ colors.

Keywords: Acyclic edge coloring; Acyclic chromatic index; κ-deletion-minimal graph; κ-minimal graph; Acyclic edge coloring conjecture

1 Introduction

All graphs considered in this paper are finite, simple and undirected. An acyclic edge coloring of a graph G is a proper edge coloring such that every cycle is colored with at least three colors. The acyclic chromatic index $\chi'_a(G)$ of a graph G is the least number of colors in an acyclic edge coloring of G. It is obvious that $\chi'_a(G) \geq \chi'(G) \geq \Delta(G)$. Fiamčík [5] stated the following conjecture in 1978, which is well known as Acyclic Edge Coloring Conjecture, and Alon et al. [2] restated it in 2001.

Conjecture 1. For any graph G, $\chi'_a(G) \leq \Delta(G) + 2$.

Alon et al. [1] proved that $\chi'_a(G) \leq 64\Delta(G)$ for any graph G by using probabilistic method. Molloy and Reed [11] improved it to $\chi'_a(G) \leq 16\Delta(G)$. Recently, Ndreca et al. [12] improved the upper bound to $\lceil 9.62(\Delta(G)-1)\rceil$, and Esperet and Parreau [4] further improved it to $4\Delta(G) - 4$ by using the so-called entropy compression method. The best known general bound is $\lceil 3.74(\Delta(G) - 1)\rceil$ due to Giotis et al. [7]. Alon et al. [2] proved that there is a constant c such that $\chi'_a(G) \leq \Delta(G) + 2$ for a graph G whenever the girth is at least $c\Delta \log \Delta$.

Regarding general planar graph G, Fiedorowicz et al. [6] proved that $\chi'_a(G) \leq 2\Delta(G) + 29$; Hou et al. [10] proved that $\chi'_a(G) \leq \max\{2\Delta(G) - 2, \Delta(G) + 22\}$. Recently, Basavaraju et al. [3] showed that $\chi'_a(G) \leq \Delta(G) + 12$, and Guan et al. [8] improved it to $\chi'_a(G) \leq \Delta(G) + 10$, and Wang et al. [14] further improved it to $\chi'_a(G) \leq \Delta(G) + 7$.

In this paper, we improve the upper bound to $\Delta(G) + 6$ by the following theorem.

Theorem 1.1. If G is a planar graph, then $\chi'_a(G) \leq \Delta(G) + 6$.

2 Preliminary

Let $\mathbb{S}$ be a multiset and x be an element in $\mathbb{S}$. The *multiplicity* $\mathrm{mul}_{\mathbb{S}}(x)$ is the number of times x appears in $\mathbb{S}$. Let $\mathbb{S}$ and $\mathbb{T}$ be two multisets. The union of $\mathbb{S}$ and $\mathbb{T}$, denoted by $\mathbb{S} \uplus \mathbb{T}$, is a multiset with $\mathrm{mul}_{\mathbb{S}\uplus\mathbb{T}}(x) = \mathrm{mul}_{\mathbb{S}}(x) + \mathrm{mul}_{\mathbb{T}}(x)$. Throughout this paper, every coloring uses colors from $[\kappa] = \{1, 2, \ldots, \kappa\}$.

We use $V(G)$, $E(G)$, $\delta(G)$ and $\Delta(G)$ to denote the vertex set, the edge set, the minimum degree and the maximum degree of a graph G, respectively. For a vertex $v \in V(G)$, $N_G(v)$ denotes the set of vertices that are adjacent to v in G and

*Corresponding author: wangtao@henu.edu.cn

$\deg_G(v)$ (or simple $\deg(v)$) to denote the degree of v in G. When G is a plane graph, we use $F(G)$ to denote its face set and $\deg_G(f)$ (or simple $\deg(f)$) to denote the degree of a face f in G. A k-, k^+-, k^--vertex (resp. face) is a vertex (resp. face) with degree k, at least k and at most k, respectively. A face $f = v_1v_2\dots v_k$ is a $(\deg(v_1), \deg(v_2), \dots, \deg(v_k))$-face.

A graph G with maximum degree at most κ is *κ-deletion-minimal* if $\chi'_a(G) > \kappa$ and $\chi'_a(H) \leq \kappa$ for every proper subgraph H of G. A graph property $\mathcal{P}$ is *deletion-closed* if $\mathcal{P}$ is closed under taking subgraphs. Analogously, we can define another type of minimal graphs by taking minors. A graph G with maximum degree at most κ is *κ-minimal* if $\chi'_a(G) > \kappa$ and $\chi'_a(H) \leq \kappa$ for every proper minor H with $\Delta(H) \leq \Delta(G)$. Obviously, every proper subgraph of a κ-minimal graph admits an acyclic edge coloring with at most κ colors, and then every κ-minimal graph is also a κ-deletion-minimal graph and all the properties of κ-deletion-minimal graphs are also true for κ-minimal graphs.

Let G be a graph and H be a subgraph of G. An acyclic edge coloring of H is a *partial acyclic edge coloring* of G. Let $\mathcal{U}_\phi(v)$ denote the set of colors which are assigned to the edges incident with v with respect to ϕ. Let $C_\phi(v) = [\kappa] \setminus \mathcal{U}_\phi(v)$ and $C_\phi(uv) = [\kappa] \setminus (\mathcal{U}_\phi(u) \cup \mathcal{U}_\phi(v))$. Let $\Upsilon_\phi(uv) = \mathcal{U}_\phi(v) \setminus \{\phi(uv)\}$ and $W_\phi(uv) = \{\, u_i \mid uu_i \in E(G)$ and $\phi(uu_i) \in \Upsilon_\phi(uv)\,\}$. Notice that $W_\phi(uv)$ may be not same with $W_\phi(vu)$. For simplicity, we will omit the subscripts if no confusion can arise.

An (α,β)-maximal dichromatic path with respect to ϕ is a maximal path whose edges are colored by α and β alternately. An (α,β,u,v)-critical path with respect to ϕ is an (α,β)-maximal dichromatic path which starts at u with color α and ends at v with color α. An (α,β,u,v)-alternating path with respect to ϕ is an (α,β)-dichromatic path starting at u with color α and ending at v with color β.

Let ϕ be a partial acyclic edge coloring of G. A color α is *candidate* for an edge e in G with respect to a partial edge coloring of G if none of the adjacent edges of e is colored with α. A candidate color α is *valid* for an edge e if assigning the color α to e does not result in any dichromatic cycle in G.

Fact 1 (Basavaraju et al. [3])**.** Given a partial acyclic edge coloring of G and two colors α,β, there exists at most one (α,β)-maximal dichromatic path containing a particular vertex v. □

Fact 2 (Basavaraju et al. [3])**.** Let G be a κ-deletion-minimal graph and uv be an edge of G. If ϕ is an acyclic edge coloring of $G-uv$, then no candidate color for uv is valid. Furthermore, if $\mathcal{U}(u)\cap\mathcal{U}(v) = \emptyset$, then $\deg(u)+\deg(v) = \kappa+2$; if $|\mathcal{U}(u) \cap \mathcal{U}(v)| = s$, then $\deg(u) + \deg(v) + \sum_{w\in W(uv)} \deg(w) \geq \kappa + 2s + 2$. □

We remind the readers that we will use these two facts frequently, so please keep these in mind and we will not refer it at every time.

3 Structural lemmas

Wang and Zhang [13] presented many structural results on κ-deletion-minimal graphs and κ-minimal graphs. In this section, we give more structural lemmas in order to prove our main result.

Lemma 1. If G is a κ-deletion-minimal graph, then G is 2-connected and $\delta(G) \geq 2$.

3.1 Local structure on the 2- or 3-vertices

Lemma 2 (Wang and Zhang [13])**.** Let G be a κ-minimal graph with $\kappa \geq \Delta(G) + 1$. If v_0 is a 2-vertex of G, then v_0 is contained in a triangle.

Lemma 3 (Wang and Zhang [13])**.** Let G be a κ-deletion-minimal graph. If v is adjacent to a 2-vertex v_0 and $N_G(v_0) = \{w, v\}$, then v is adjacent to at least $\kappa - \deg(w) + 1$ vertices with degree at least $\kappa - \deg(v) + 2$. Moreover,

(A) if $\kappa \geq \deg(v) + 1$ and $wv \in E(G)$, then v is adjacent to at least $\kappa - \deg(w) + 2$ vertices with degree at least $\kappa - \deg(v) + 2$, and $\deg(v) \geq \kappa - \deg(w) + 3$;

(B) if $\kappa \geq \Delta(G) + 2$ and v is adjacent to precisely $\kappa - \Delta(G) + 1$ vertices with degree at least $\kappa - \Delta(G) + 2$, then v is adjacent to at most $\deg(v) + \Delta(G) - \kappa - 3$ vertices with degree two and $\deg(v) \geq \kappa - \Delta(G) + 4$.

Lemma 4 (Wang and Zhang [13])**.** Let G be a κ-deletion-minimal graph with $\kappa \geq \Delta(G) + 2$. If v_0 is a 2-vertex, then every neighbor of v_0 has degree at least $\kappa - \Delta(G) + 4$.

Lemma 5 (Hou et al. [9]). Let G be a κ-deletion-minimal graph with $\kappa \geq \Delta(G) + 2$. If v is a 3-vertex, then every neighbor of v is a $(\kappa - \Delta(G) + 2)^+$-vertex.

Lemma 6 (Wang and Zhang [13]). Let G be a κ-minimal graph with $\kappa \geq \Delta(G) + 2$. If v is a 3-vertex in G, then every neighbor of v is a $(\kappa - \Delta(G) + 3)^+$-vertex.

Lemma 7 (Wang and Zhang [13]). Let G be a κ-deletion-minimal graph with $\kappa \geq \Delta(G) + 2$, and let w_0 be a 3-vertex with $N_G(w_0) = \{w, w_1, w_2\}$, and $\deg(w) = \kappa - \Delta(G) + 3$. If $ww_1, ww_2 \in E(G)$, then $\deg(w_1) = \deg(w_2) = \Delta(G)$ and w is adjacent to precisely one vertex (namely w_0) with degree less than $\Delta(G) - 1$.

Lemma 8. Let G be a κ-deletion-minimal graph with maximum degree Δ, and let w_0 be a 3-vertex with $N_G(w_0) = \{w, w_1, w_2\}$. If $\deg_G(w) = \kappa - \Delta + 4 = \ell$ with $8 \leq \ell \leq 10$ and $N_G(w) = \{w_0, w_1, w_2, \ldots, w_{\ell-1}\}$, then there exists no 4-set $X^* \subseteq \{w_1, w_2, \ldots, w_{\ell-1}\}$ satisfying the following four conditions: (1) every vertex in X^* is a 5^--vertex; (2) the degree-sum of vertices in X^* is at most $\kappa - \Delta + 9$; (3) the degree-sum of any two vertices in X^* is at most Δ; (4) X^* has at least two 4^--vertices.

Proof. Suppose to the contrary that there exists a 4-set X^* satisfying all the four conditions. Let X be the subscripts of vertices in X^*. Since G is κ-deletion-minimal, the graph $G - ww_0$ has an acyclic edge coloring ϕ with $\phi(ww_i) = i$ for $i \in \{1, \ldots, \ell - 1\}$. The fact that $\deg_G(w) + \deg_G(w_0) \leq \Delta + 3 < \kappa + 2$ and Fact 2 imply that $\mathcal{U}(w) \cap \mathcal{U}(w_0) \neq \emptyset$.

Case 1. $|\mathcal{U}(w) \cap \mathcal{U}(w_0)| = 1$.

It follows that $|C(ww_0)| = \Delta - 4$.

Subcase 1.1. The common color is on ww_1 or ww_2.

Without loss of generality, we may assume that w_0w_1 is colored with ℓ and w_0w_2 is colored with 1. Note that there exists a $(1, \alpha, w, w_0)$-critical path for every $\alpha \in \{\ell+1, \ldots, \kappa\}$, so we have that $\{\ell+1, \ldots, \kappa\} \subseteq \mathcal{U}(w_1) \cap \mathcal{U}(w_2)$. Notice that the set $\{1, \ldots, \ell\} \setminus (\mathcal{U}(w_1) \cup \mathcal{U}(w_2))$ is nonempty. Now, reassigning ℓ to ww_0 and a color in $\{1, \ldots, \ell\} \setminus (\mathcal{U}(w_1) \cup \mathcal{U}(w_2))$ to w_0w_1 results in an acyclic edge coloring of G, a contradiction.

Subcase 1.2. The common color is not on ww_1 and ww_2.

Without loss of generality, we may assume that w_0w_1 is colored with ℓ and w_0w_2 is colored with 3. There exists a $(3, \alpha, w, w_0)$-critical path for $\alpha \in \{\ell + 1, \ldots, \kappa\}$. It follows that $\{\ell + 1, \ldots, \kappa\} \subseteq \Upsilon(ww_3) \cap \Upsilon(w_0w_2)$ and $\deg_G(w_3) \geq \Delta - 3 \geq 5$.

If $1 \notin \mathcal{U}(w_2)$, then reassigning 1 to w_0w_2 will take us back to Case 1.1. Hence, we have that $1 \in \Upsilon(w_0w_2)$ and $\deg_G(w_2) \geq \Delta - 1 \geq 7$. By Lemma 5, we have that $\deg_G(w_1) \geq \kappa - \Delta + 2 \geq 6$.

Note that w_1, w_2 and w_3 are 5^+-vertices, there exists a 4^--vertex w_x with $x \in X \setminus \mathcal{U}(w_2)$. If $\ell \notin \mathcal{U}(w_2)$, then reassigning the color x to w_0w_2 results in a new acyclic edge coloring σ of $G - ww_0$, and then $C_\sigma(ww_0) = \{\ell+1, \ldots, \kappa\} \subseteq \Upsilon(ww_x)$ and $\deg_G(w_x) \geq \Delta - 3 \geq 5$, which contradicts that w_x is a 4^--vertex. Hence, $\Upsilon(w_0w_2) = \{1, 2\} \cup \{\ell, \ldots, \kappa\}$ and $\deg_G(w_2) = \Delta$, which implies that $X \cap \Upsilon(w_0w_2) = \emptyset$.

Claim 1. There exists a $(3, \ell, w, w_2)$-alternating path.

Proof. Suppose to the contrary that there exists no $(3, \ell, w, w_2)$-alternating path. We can revise ϕ by assigning ℓ to ww_0 and erase the color from w_0w_1, and obtain an acyclic edge coloring of $G - w_0w_1$. If some color $\alpha \in \{\ell+1, \ldots, \kappa\}$ is absent in $\mathcal{U}_\phi(w_1)$, then we can further assign α to w_0w_1, since there exists a $(3, \alpha, w, w_0)$-critical path with respect to ϕ. If some color $\alpha \in \{4, \ldots, \ell - 1\}$ is absent in $\mathcal{U}_\phi(w_1)$, then we can further assign α to w_0w_1. Hence, $\mathcal{U}_\phi(w_1) \supseteq \{1\} \cup \{4, \ldots, \kappa\}$ and $\deg_G(w_1) \geq \kappa - 2 > \Delta(G)$, a contradiction. □

Therefore, $\{\ell, \ldots, \kappa\} \subseteq \Upsilon(ww_3)$ and $\deg_G(w_3) \geq \Delta - 2 \geq 6$, which implies that $X \cap \mathcal{U}(w_2) = \emptyset$.

There exists a (ℓ, m, w_0, w_2)-critical path for every $m \in X$; otherwise, reassigning m to w_0w_2 results in another new acyclic edge coloring ϕ_m of $G - ww_0$, by the above arguments, $\{\ell, \ldots, \kappa\} \subseteq \Upsilon(ww_m)$ and $\deg_G(w_m) \geq \Delta - 2 \geq 6$, a contradiction. Thus, we have that $X \subseteq \Upsilon(w_0w_1)$. By symmetry, we may assume that $\{4, 5, 6, 7\} = X \subseteq \Upsilon(w_0w_1)$.

Suppose that $\{3, 8, 9, \ldots, \ell - 1\} \nsubseteq \mathcal{U}(w_1)$, say λ is a such color. There exists a $(\lambda, \alpha, w, w_2)$-alternating path for $\ell + 1 \leq \alpha \leq \kappa$; otherwise, reassigning λ to w_0w_2 (if $\lambda = 3$ there is no change to w_0w_2) and α to ww_0 results in an acyclic edge coloring of G. Similar to Claim 1, there exists a (λ, ℓ, w, w_2)-alternating path. Reassigning λ to w_0w_1 and 4 to w_0w_2 results in a new acyclic edge coloring φ of $G - ww_0$. Since there is no $(\lambda, \alpha, w, w_0)$-critical path with respect to

φ, thus there exists a $(4, \alpha, w_0, w)$-critical path with respect to φ for $\alpha \in \{\ell, \dots, \kappa\}$, and then $\{\ell, \dots, \kappa\} \subseteq \Upsilon(ww_4)$, which contradicts the fact that w_4 is a 5^--vertex. Hence, we have that $\{1\} \cup \{3, 4, \dots, \ell\} \subseteq \mathcal{U}(w_1)$.

Let φ_m be obtained from ϕ by reassigning m to ww_0 and erasing the color on ww_m, where $m \in \{4, 5, 6, 7\}$. Note that φ_m is an acyclic edge coloring of $G - ww_m$ for $m \in \{4, 5, 6, 7\}$. By Fact 2, we have that $|\Upsilon(ww_m) \cap \{1, 2, \dots, \ell - 1\}| \geq 1$ for $m \in \{4, 5, 6, 7\}$.

Let α be an arbitrary color in $\{\ell, \dots, \kappa\} \setminus (\Upsilon(w_0w_1) \cup \Upsilon(ww_4) \cup \Upsilon(ww_5) \cup \Upsilon(ww_6) \cup \Upsilon(ww_7))$. Since there exists neither $(1, \alpha, w, w_x)$-critical path nor $(3, \alpha, w, w_x)$-critical path (with respect to φ_x) for every $x \in X$, thus there exists a $(\lambda_x, \alpha, w, w_x)$-critical path (with respect to φ_x), where $\lambda_x \in \{2, 8, 9, \dots, \ell - 1\}$. Moreover, there exists $(\lambda, \alpha, w, w_{x_1})$- and $(\lambda, \alpha, w, w_{x_2})$-critical path for some $\lambda \in \{2, 8, 9, \dots, \ell - 1\}$ since $|X| > |\{2, 8, 9, \dots, \ell - 1\}|$, but this contradicts Fact 1.

So we may assume that $\alpha \in \Upsilon(ww_4) \cup \Upsilon(ww_5) \cup \Upsilon(ww_6) \cup \Upsilon(ww_7)$ for every $\alpha \in \{\ell, \dots, \kappa\} \setminus \Upsilon(w_0w_1)$.

$$\begin{aligned}
\kappa - \Delta + 9 \geq& \deg_G(w_4) + \deg_G(w_5) + \deg_G(w_6) + \deg_G(w_7) \\
\geq& |\{\ell, \dots, \kappa\} \setminus \Upsilon(w_0w_1)| + 4 + \sum_{t=4}^{7} |\Upsilon(ww_t) \cap \{1, \dots, \ell - 1\}| \\
\geq& (\kappa - \Delta) + 4 + (1 + 1 + 1 + 1) \\
=& \kappa - \Delta + 8.
\end{aligned}$$

By symmetry, we may assume that $|\Upsilon(ww_4) \cap \{1, \dots, \ell-1\}| = |\Upsilon(ww_5) \cap \{1, \dots, \ell-1\}| = |\Upsilon(ww_6) \cap \{1, \dots, \ell-1\}| = 1$. Let $\Upsilon(ww_4) \cap \{1, \dots, \ell-1\} = \{\mu_1\}$, $\Upsilon(ww_5) \cap \{1, \dots, \ell-1\} = \{\mu_2\}$ and $\Upsilon(ww_6) \cap \{1, \dots, \ell-1\} = \{\mu_3\}$. If $\mu_1 = \mu_2 = \mu$, then there exists a (μ, α, w, w_4)- and (μ, α, w, w_5)-critical path, where $\alpha \in \{\ell, \dots, \kappa\} \setminus (\Upsilon(ww_4) \cup \Upsilon(ww_5))$, which contradicts Fact 1. Thus μ_1, μ_2, μ_3 are distinct.

If $\mu_1 \in \{4, 5, 6, 7\}$, then every color $\alpha \in \{\ell, \dots, \kappa\} \setminus (\Upsilon(ww_4) \cup \Upsilon(ww_{\mu_1}))$ is valid for ww_4 with respect to φ_4; note that $\{\ell, \dots, \kappa\} \setminus (\Upsilon(ww_4) \cup \Upsilon(ww_{\mu_1}))$ is a nonempty set. By symmetry, we may assume that $\{\mu_1, \mu_2, \mu_3\} \cap \{4, 5, 6, 7\} = \emptyset$.

Since μ_1, μ_2, μ_3 are distinct, we may assume that $\mu_1 \neq 2$. If $2 \notin \Upsilon(w_0w_1)$, then reassigning 2 to w_0w_1 and 4 to w_0w_2 results in a new acyclic edge coloring φ^* of $G - ww_0$. For every color $\beta \in \{\ell, \dots, \kappa\} \setminus \Upsilon(w_0w_1)$, there exists no $(2, \beta, w, w_0)$-critical path with respect to φ^*, thus there exists a $(4, \beta, w, w_0)$-critical path with respect to φ^*, and then $\{\ell, \dots, \kappa\} \setminus \Upsilon(w_0w_1) \subseteq \Upsilon(ww_4)$ and $\deg_G(w_4) \geq |\{\ell, \dots, \kappa\} \setminus \Upsilon(w_0w_1)| + 2 \geq 6$, which contradicts the degree of w_4.

Hence, we have that $\{1, \dots, \ell - 1\} \subseteq \Upsilon(w_0w_1)$ and $|\{\ell, \dots, \kappa\} \setminus \Upsilon(w_0w_1)| \geq \kappa - \Delta + 1$. By similar arguments as above, we can prove that $\Upsilon(ww_7) \cap \{1, \dots, \ell - 1\} = \{\mu_4\}$ and $\mu_1, \mu_2, \mu_3, \mu_4$ are distinct. Moreover, we can also conclude that $\{\mu_1, \mu_2, \mu_3, \mu_4\} \cap \{4, 5, 6, 7\} = \emptyset$.

Suppose that $\mu_1 = 3$. Since there exists no $(3, \alpha, w, w_4)$-critical path with respect to φ_4, where $\alpha \in \{\ell + 1, \dots, \kappa\}$, thus $\{\ell + 1, \dots, \kappa\} \subseteq \Upsilon(ww_4)$, a contradiction. So, by symmetry, we may assume that $\{\mu_1, \mu_2, \mu_3, \mu_4\} = \{1, 2, 8, 9\}$.

By symmetry, we assume that $\mu_1 = 1$. Note that there exists no $(1, \alpha, w, w_4)$-critical path (with respect to φ_4) for every $\alpha \in \{\ell, \dots, \kappa\} \setminus \Upsilon(w_0w_1)$, thus $\{\ell, \dots, \kappa\} \setminus \Upsilon(w_0w_1) \subseteq \Upsilon(ww_4)$; otherwise, reassigning 4 to ww_0 and a color α to ww_4 results in an acyclic edge coloring. Now, we have that $\deg_G(w_4) \geq 2 + |\{\ell, \dots, \kappa\} \setminus \Upsilon(w_0w_1)| \geq 6$, a contradiction.

Case 2. $\mathcal{U}(w) \cap \mathcal{U}(w_0) = \{\lambda_1, \lambda_2\}$, $\phi(w_0w_1) = \lambda_1$ and $\phi(w_0w_2) = \lambda_2$.

If follows that $|C(ww_0)| = \Delta - 3$. First of all, we show the following claim:

$(*)$ $C(ww_0) = \{\ell, \dots, \kappa\} \subseteq \mathcal{U}(w_1) \cap \mathcal{U}(w_2)$.

By contradiction and symmetry, assume that there exists a color ζ in $\{\ell, \dots, \kappa\} \setminus \mathcal{U}(w_1)$. Clearly, there exists a $(\lambda_2, \zeta, w_0, w)$-critical path, and then there exists no $(\lambda_2, \zeta, w_0, w_1)$-critical path. Now, reassigning ζ to w_0w_1 will take us back to Case 1. Hence, we have $\{\ell, \dots, \kappa\} \subseteq \mathcal{U}(w_1)$; similarly, we have $\{\ell, \dots, \kappa\} \subseteq \mathcal{U}(w_2)$. This completes the proof of $(*)$.

Note that w_1 and w_2 have degree at least $\Delta - 1 \geq 7$, this implies that $\{1, 2\} \cap X = \emptyset$ and $|X \cap \Upsilon(w_0w_1)| \leq 1$ and $|X \cap \Upsilon(w_0w_2)| \leq 1$. Let $\{p, q\} \subseteq X \setminus (\Upsilon(w_0w_1) \cup \Upsilon(w_0w_2))$. Reassigning p to w_0w_1 and q to w_0w_2 results in a new acyclic edge coloring ψ of $G - ww_0$. Hence, we have that $C_\psi(ww_0) \subseteq \Upsilon(ww_p) \cup \Upsilon(ww_q)$, and then $\deg_G(w_p) + \deg_G(w_q) \geq (\Delta - 3) + 2 + 2 \geq \Delta + 1$, which is a contradiction. □

3.2 Local structure on the 4-vertices

Lemma 9. Let G be a κ-deletion-minimal graph with maximum degree Δ and $\kappa \geq \Delta + 2$, and let w_0 be a 4-vertex with $N_G(w_0) = \{w, v_1, v_2, v_3\}$.

(a) If $\deg_G(w) \le \kappa - \Delta$, then

$$\sum_{x \in N_G(w_0)} \deg_G(x) \ge 2\kappa - \deg_G(w_0) + 8 = 2\kappa + 4. \tag{1}$$

(b) If $\deg_G(w) \le \kappa - \Delta + 1$ and ww_0 is contained in two triangles ww_1w_0 and ww_2w_0, then

$$\sum_{x \in N_G(w_0)} \deg_G(x) \ge 2\kappa - \deg_G(w_0) + 9 = 2\kappa + 5. \tag{2}$$

Furthermore, if the equality holds in (2), then all the other neighbors of w are 6^+-vertices.

Proof. We may assume that

(∗) The graph $G - ww_0$ admits an acyclic edge coloring ϕ such that the number of common colors at w and w_0 is minimum.

Here, (a) and (b) will be proved together, so we may assume that $\deg_G(w) \le \kappa - \Delta + 1$. Since $\deg_G(w) + \deg_G(w_0) \le \kappa - \Delta + 5 < \kappa + 2$, we have that $|\Upsilon(ww_0) \cap \Upsilon(w_0w)| = m \ge 1$. It follows that $|C(ww_0)| = \kappa - (\deg_G(w) + \deg_G(w_0) - m - 2) \ge \Delta - 2$. Without loss of generality, let $N_G(w) = \{w_0, w_1, w_2, \dots\}$ and $\phi(ww_i) = i$ for $1 \le i \le \deg_G(w) - 1$. Let $\mathbb{S} = \Upsilon(w_0v_1) \uplus \Upsilon(w_0v_2) \uplus \Upsilon(w_0v_3)$.

Claim 1. For every color θ in $C(ww_0)$, there exists a $(\lambda, \theta, w_0, w)$-critical path for some $\lambda \in \Upsilon(ww_0) \cap \Upsilon(w_0w)$. Consequently, every color in $C(ww_0)$ appears in $\mathbb{S}$.

Case 1. $\mathcal{U}(w) \cap \mathcal{U}(w_0) = \{\lambda\}$.

It follows that $|C(ww_0)| = \kappa - (\deg_G(w) + \deg_G(w_0) - 3)$.

(a) Suppose that $\deg_G(w) + \deg_G(w_0) \le \kappa - \Delta + 4$. It follows that $|C(ww_0)| \ge \Delta - 1$. Without loss of generality, let $\phi(w_0v_1) = 1, \phi(w_0v_2) = \kappa - \Delta$ and $\phi(w_0v_3) = \kappa - \Delta + 1$. By Claim 1, there exists a $(1, \theta, w_0, w)$-critical path for every θ in $C(ww_0)$. Hence, we have that $\deg_G(w) = \kappa - \Delta$ and $\deg_G(v_1) = \deg_G(w_1) = \Delta$ and $\Upsilon(w_0v_1) = \Upsilon(ww_1) = \{\kappa - \Delta + 2, \dots, \kappa\}$. Notice that $\deg_G(w) = \kappa - \Delta \ge 3$ results from Lemma 4. Reassigning $\kappa - \Delta$, 1 and 2 to ww_1, ww_0 and w_0v_1 respectively, and we obtain an acyclic edge coloring of G, a contradiction.

(b) Suppose that $\deg_G(w) + \deg_G(w_0) = \kappa - \Delta + 5$ and ww_0 is contained in two triangles ww_1w_0 and ww_2w_0 ($w_1 = v_1$ and $w_2 = v_2$).

Subcase 1.1. The common color λ does not appear on w_0v_3, but it appears on ww_1 or ww_2.

By symmetry, assume that $\phi(w_0w_1) = 2$, $\phi(w_0v_2) = \kappa - \Delta + 1$, $\phi(w_0v_3) = \kappa - \Delta + 2$. By Claim 1, we have that $\{\kappa - \Delta + 3, \dots, \kappa\} \subseteq \Upsilon(w_0w_1) \cap \Upsilon(ww_2)$ and $\deg_G(w_1) = \deg_G(w_2) = \Delta$. Now, reassigning $\kappa - \Delta + 1$ to w_0w and reassigning 3 to w_0w_2 results in an acyclic edge coloring of G, a contradiction.

Subcase 1.2. The common color λ does not appear on w_0v_3 and it does not appear on ww_1 or ww_2 either.

By symmetry, assume that $\phi(w_0w_1) = 3$, $\phi(w_0w_2) = \kappa - \Delta + 1$, $\phi(w_0v_3) = \kappa - \Delta + 2$. By Claim 1, we have that $\{\kappa - \Delta + 3, \dots, \kappa\} \subseteq \Upsilon(w_0w_1) \cap \Upsilon(ww_3)$, $\deg_G(w_1) = \Delta$ and $\deg_G(w_3) \ge \Delta - 1$. Reassigning 2 to w_0w_1 will take us back to Subcase 1.1.

Subcase 1.3. The common color λ appears on w_0v_3 and it also appears on ww_1 or ww_2.

By symmetry, assume that $\phi(w_0w_1) = \kappa - \Delta + 1$, $\phi(w_0w_2) = \kappa - \Delta + 2$, $\phi(w_0v_3) = 2$. By Claim 1, we have that $\{\kappa - \Delta + 3, \dots, \kappa\} \subseteq \Upsilon(ww_2) \cap \Upsilon(w_0v_3)$, $\deg_G(w_2) = \Delta$ and $\deg_G(v_3) \ge \Delta - 1$. Now, reassigning $\kappa - \Delta + 1$ to ww_2 will take us back to Subcase 1.1.

Subcase 1.4. The common color λ appears on w_0v_3, but it does not appear on ww_1 or ww_2.

By symmetry, assume that $\phi(w_0w_1) = \kappa - \Delta + 1$, $\phi(w_0w_2) = \kappa - \Delta + 2$, $\phi(w_0v_3) = 3$. By Claim 1, we have that $\{\kappa - \Delta + 3, \dots, \kappa\} \subseteq \Upsilon(ww_3) \cap \Upsilon(w_0v_3)$, $\deg_G(w_3) \ge \Delta - 1$ and $\deg_G(v_3) \ge \Delta - 1$. If $\{2, \kappa - \Delta + 1\} \cap \Upsilon(w_0v_3) = \emptyset$, then reassigning 2 to w_0v_3 will take us back to Subcase 1.3. So we may assume that $\{2, \kappa - \Delta + 1\} \cap \Upsilon(w_0v_3) \ne \emptyset$. But we can still reassign 1 to w_0v_3 and go back to Subcase 1.3.

Case 2. $\mathcal{U}(w) \cap \mathcal{U}(w_0) = \{\lambda_1, \dots, \lambda_m\}$ and $m \ge 2$.

Let $\mathcal{A}(v_1) = C(ww_0) \setminus \Upsilon(w_0v_1) = \{\alpha_1, \alpha_2, \dots\}$, $\mathcal{A}(v_2) = C(ww_0) \setminus \Upsilon(w_0v_2) = \{\beta_1, \beta_2, \dots\}$ and $\mathcal{A}(v_3) = C(ww_0) \setminus \Upsilon(w_0v_3)$.

Claim 2. $\mathcal{A}(v_1), \mathcal{A}(v_2), \mathcal{A}(v_3) \neq \emptyset$.

Proof. Suppose to the contrary that $\mathcal{A}(v_*) = \emptyset$. It follows that $\Delta - 1 \geq |\Upsilon(w_0v_*)| \geq |C(ww_0)| = \kappa - (\deg_G(w) + \deg_G(w_0) - m - 2) \geq \kappa - (\kappa - \Delta + 5 - 2 - 2) = \Delta - 1$, thus $\deg_G(w) + \deg_G(w_0) = \kappa - \Delta + 5$, $m = 2$ and $\Upsilon(w_0v_*) = C(ww_0)$ with $|\Upsilon(w_0v_*)| = \Delta - 1$. This implies that the graph G satisfies the condition (b) with $v_* = v_3$ (assume that $w_1 = v_1$ and $w_2 = v_2$). We may assume that $\mathcal{U}(w_0) = \{\lambda_1, \lambda_2, \kappa - \Delta + 1\}$.

If the color on w_0w_1 is λ_1 and the color on w_0w_2 is λ_2, then reassigning α_1, β_1 and λ_2 to ww_0, w_0w_2 and w_0v_3, respectively, yields an acyclic edge coloring of G.

But if the color on w_0w_1 is $\kappa - \Delta + 1$ and the color on w_0w_2 is λ_2, then reassigning 2 to w_0v_3 and β_1 to ww_0 results in an acyclic edge coloring of G. □

Claim 3. Every color in $C(ww_0)$ appears at least twice in $\mathbb{S}$.

Proof. Suppose that there exists a color α in $C(ww_0)$ appearing only once in $\mathbb{S}$, say $\alpha \in \Upsilon(w_0v_1)$. Without loss of generality, assume that $\phi(w_0v_1) = \lambda_1$ and $\phi(w_0v_2) = \lambda_2$. By Claim 1, there exists a $(\lambda_1, \alpha, w_0, w)$-critical path. Reassigning α to w_0v_2 results in a new acyclic edge coloring ϕ^* of $G - ww_0$ with $|\mathcal{U}_{\phi^*}(w) \cap \mathcal{U}_{\phi^*}(w_0)| < |\mathcal{U}(w) \cap \mathcal{U}(w_0)|$, which contradicts the assumption $(*)$. □

Let $X = \{\, \alpha \mid \alpha \in C(ww_0) \text{ and } \mathrm{mul}_{\mathbb{S}}(\alpha) = 3 \,\}$.

$$
\begin{aligned}
&\sum_{x \in N_G(w_0)} \deg_G(x) \\
&= \deg_G(w_0) + \deg_G(w) - 1 + \sum_{\alpha \in [\kappa]} \mathrm{mul}_{\mathbb{S}}(\alpha) \\
&= \deg_G(w_0) + \deg_G(w) - 1 + \sum_{\alpha \in C(ww_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) + \sum_{\alpha \in \mathcal{U}(w) \cup \mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) \\
&= \deg_G(w_0) + \deg_G(w) - 1 + 2|C(ww_0)| + |X| + \sum_{\alpha \in \mathcal{U}(w) \cup \mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) \\
&= \deg_G(w_0) + \deg_G(w) - 1 + 2(\kappa - (\deg_G(w) + \deg_G(w_0) - 2 - m)) + |X| + \sum_{\alpha \in \mathcal{U}(w) \cup \mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) \\
&= 2\kappa - \deg_G(w_0) - \deg_G(w) + 2m + 3 + |X| + \sum_{\alpha \in \mathcal{U}(w) \cup \mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha)
\end{aligned}
$$

It is sufficient to prove that

$$
\sum_{\alpha \in \mathcal{U}(w) \cup \mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) + |X| \geq \begin{cases} \deg_G(w) - 2m + 5, & \text{if } \deg_G(w) \leq \kappa - \Delta; \quad (3) \\ \deg_G(w) - 2m + 6, & \text{if } \deg_G(w) \leq \kappa - \Delta + 1 \text{ and } ww_0 \text{ is contained in two triangles.} \quad (4) \end{cases}
$$

Subcase 2.1. $\mathcal{U}(w) \cap \mathcal{U}(w_0) = \{\lambda_1, \lambda_2\}$.

Claim 4. Every color in $\mathcal{U}(w)$ is in $\mathbb{S}$.

Proof. Assume that w_0v_1 is colored with λ_1 and w_0v_2 is colored with λ_2. Notice that $C(ww_0) \subseteq \Upsilon(w_0v_1) \cup \Upsilon(w_0v_2)$ and $\mathcal{A}(v_1) \cap \mathcal{A}(v_2) = \emptyset$. By Claim 2, we have that $\mathcal{A}(v_1), \mathcal{A}(v_2), \mathcal{A}(v_3) \neq \emptyset$. If $\lambda_1 \notin \mathbb{S}$, then reassigning β_1, α_1 and λ_1 to w_0w, w_0v_1 and w_0v_3 respectively, results in an acyclic edge coloring of G, a contradiction. Thus, we have that $\lambda_1 \in \mathbb{S}$. Similarly, we can prove that $\lambda_2 \in \mathbb{S}$. Let τ be an arbitrary color in $\mathcal{U}(w) \setminus (\mathbb{S} \cup \{\lambda_1, \lambda_2\})$. Let σ be obtained from ϕ by reassigning τ to w_0v_1. It is obvious that σ is an acyclic edge coloring of $G - ww_0$. So we can obtain a similar contradiction by replacing ϕ with σ. □

Claim 5. The color in $\mathcal{U}(w_0) \setminus \{\lambda_1, \lambda_2\}$ appears at least twice in $\mathbb{S}$.

Proof. Suppose that λ_1, λ_2 and λ^* are on the edges w_0v_1, w_0v_2 and w_0v_3, respectively. There exists a $(\lambda^*, \alpha_1, w_0, v_1)$-critical path; otherwise, reassigning α_1 to w_0v_1 will take us back to Case 1. Hence, we have $\lambda^* \in \Upsilon(w_0v_1)$. Similarly, there exists a $(\lambda^*, \beta_1, w_0, v_2)$-critical path and $\lambda^* \in \Upsilon(w_0v_2)$. Therefore, the color λ^* appears exactly twice in $\mathbb{S}$. □

Now, we have

$$\sum_{\alpha \in \mathcal{U}(w) \cup \mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) + |X| \geq |\mathcal{U}(w)| + 2 + |X| = \deg_G(w) + 1 + |X|.$$

So conclusion (a) holds. Now, suppose that $\deg_G(w) + \deg_G(w_0) \leq \kappa - \Delta + 5$ and ww_0 is contained in two triangles ww_0w_1 and ww_0w_2 ($w_1 = v_1$ and $w_2 = v_2$).

Subcase 2.1.1. The two common colors λ_1 and λ_2 are on w_1w and w_1w_0.

There exists a $(\lambda_1, \alpha, w_0, w)$- or $(\lambda_2, \alpha, w_0, w)$-critical path for $\alpha \in C(ww_0)$. Hence, we have that $C(ww_0) \subseteq \mathcal{U}(w_1)$, and thus $\deg_G(w_1) \geq |C(ww_0)| + |\{\lambda_1, \lambda_2\}| \geq \Delta + 1$, a contradiction.

Subcase 2.1.2. The two common colors λ_1 and λ_2 are on w_2w and w_2w_0.

This is similar with Subcase 2.1.1.

Subcase 2.1.3. $\{\lambda_1, \lambda_2\} \cap \{1, 2\} = \{\lambda_1\}$ and λ_1 appears on w_0w_1 or w_0w_2.

Without loss of generality, assume that $\phi(w_0w_1) = \kappa - \Delta + 1$, $\phi(w_0w_2) = 1$, $\phi(w_0v_3) = 3$. If $2 \notin \Upsilon(w_0w_1) \cup \Upsilon(w_0v_3)$, then reassigning 2 to w_0v_3 will take us back to Subcase 2.1.2. Hence, $2 \in \Upsilon(w_0w_1) \cup \Upsilon(w_0v_3)$ and 2 appears at least twice in $\mathbb{S}$. Therefore, we have

$$\sum_{\alpha \in \mathcal{U}(w) \cup \mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) + |X| \geq |\mathcal{U}(w)| + 2 + |X| + |\{2\}| \geq \deg_G(w) + 2.$$

Suppose that

$$\sum_{x \in N_G(w_0)} \deg_G(x) = 2\kappa - \deg_G(w_0) + 9.$$

It follows that

$$\sum_{\alpha \in \mathcal{U}(w) \cup \mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) + |X| = |\mathcal{U}(w)| + 2 + |X| + |\{2\}| = \deg_G(w) + 2,$$

and every color in $\mathcal{U}(w) \setminus \{2\}$ appears only once in $\mathbb{S}$.

There exists a $(3, \kappa - \Delta + 1, w_0, w)$-critical path, otherwise, reassigning $\kappa - \Delta + 1$ to w_0w and α_1 to w_0w_1 results in an acyclic edge coloring of G, a contradiction. By Claim 5, we have that $\kappa - \Delta + 1 \in \Upsilon(w_0w_2) \cap \Upsilon(w_0v_3)$. And by Claim 4, we have that $3 \in \Upsilon(w_0w_1) \cup \Upsilon(w_0w_2)$. Since $|C(ww_0)| \geq \Delta - 1$ and $\{1, 2, 3, \kappa - \Delta + 1\} \subseteq \Upsilon(w_0w_1) \cup \Upsilon(w_0w_2)$, this implies that $|\mathcal{A}(w_1)| + |\mathcal{A}(w_2)| \geq 4$. There exists no $(1, \alpha, w, w_0)$-critical path for every $\alpha \in \mathcal{A}(w_1) \cup \mathcal{A}(w_2)$, thus there exists a $(3, \alpha, w, w_0)$-critical path, and then $\mathcal{A}(w_1) \cup \mathcal{A}(w_2) \subseteq \Upsilon(ww_3)$. Hence, $\deg_G(w_3) \geq |\mathcal{A}(w_1)| + |\mathcal{A}(w_2)| + |\{3, \kappa - \Delta + 1\}| \geq 6$.

Suppose that $4 \notin \Upsilon(w_0v_3)$ and there exists no $(\kappa - \Delta + 1, 4, w_0, v_3)$-critical path. Reassigning 4 to w_0v_3 results in a new acyclic edge coloring ϱ_1 of $G - ww_0$. Similarly, we can prove $\deg_G(w_4) \geq 6$ by replacing ϕ with ϱ_1.

Suppose that $4 \in \Upsilon(w_0v_3)$. This implies that $\{1, 2, 4, \kappa - \Delta + 1\} \subseteq \Upsilon(w_0w_1) \cup \Upsilon(w_0v_3)$ and $|\mathcal{A}(w_1)| + |\mathcal{A}(v_3)| \geq 4$. Reassigning 4 to w_0w_2 and reassigning 1 to w_0v_3 results in another acyclic edge coloring π of $G - ww_0$. Hence, there exists a $(4, \alpha, w_0, w)$-critical path with respect to π for $\alpha \in \mathcal{A}(w_1) \cup \mathcal{A}(v_3)$, and then $\mathcal{A}(w_1) \cup \mathcal{A}(v_3) \subseteq \Upsilon(ww_4)$. Similarly as above, there exists a $(4, \kappa - \Delta + 1, w_0, w)$-critical path with respect to π. Hence, $\deg_G(w_4) \geq |\mathcal{A}(w_1)| + |\mathcal{A}(v_3)| + |\{4, \kappa - \Delta + 1\}| \geq 6$.

Suppose that there exists a $(\kappa - \Delta + 1, 4, w_0, v_3)$-critical path and $4 \in \Upsilon(w_0w_1)$. This implies that $\{1, 2, 4, \kappa - \Delta + 1\} \subseteq \Upsilon(w_0w_1) \cup \Upsilon(w_0v_3)$ and $|\mathcal{A}(w_1)| + |\mathcal{A}(v_3)| \geq 4$. Reassigning 4 to w_0w_2 and reassigning 1 to w_0v_3 results in another acyclic edge coloring ϱ_2 of $G - ww_0$. Similarly as above, we can prove that $\deg_G(w_4) \geq 6$.

In one word, the degree of w_4 is at least six. By symmetry, we have that $\deg_G(w_i) \geq 6$ for $4 \leq i \leq \deg_G(w) - 1$.

Subcase 2.1.4. $\{\lambda_1, \lambda_2\} \cap \{1, 2\} = \{\lambda_1\}$ and λ_1 appears on w_0v_3.

Without loss of generality, assume that $\phi(w_0w_1) = \kappa - \Delta + 1$, $\phi(w_0w_2) = 3$, $\phi(w_0v_3) = 1$. If $2 \notin \Upsilon(w_0w_1) \cup \Upsilon(w_0v_3)$, then reassigning 2 to w_0w_1 and reassigning β_1 to w_0w_2 will take us back to Subcase 2.1.1. Hence, $2 \in \Upsilon(w_0w_1) \cup \Upsilon(w_0v_3)$ and 2 appears at least twice in $\mathbb{S}$. Therefore, we have

$$\sum_{\alpha\in\mathcal{U}(w)\cup\mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) + |X| \geq |\mathcal{U}(w)| + 2 + |X| + |\{2\}| \geq \deg_G(w) + 2.$$

Suppose that

$$\sum_{x\in N_G(w_0)} \deg_G(x) = 2\kappa - \deg_G(w_0) + 9.$$

It follows that

$$\sum_{\alpha\in\mathcal{U}(w)\cup\mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) + |X| = |\mathcal{U}(w)| + 2 + |X| + |\{2\}| = \deg_G(w) + 2,$$

and every color in $\mathcal{U}(w) \setminus \{2\}$ appears only once in $\mathbb{S}$.

There exists a $(3, \kappa-\Delta+1, w_0, w)$-critical path, otherwise, reassigning $\kappa-\Delta+1$ to w_0w and α_1 to w_0w_1 results in an acyclic edge coloring of G, a contradiction. Since $|C(ww_0)| \geq \Delta - 1$ and $\{1, 2, 3, \kappa-\Delta+1\} \subseteq \Upsilon(w_0w_1) \cup \Upsilon(w_0v_3)$, this implies that $|\mathcal{A}(w_1)|+|\mathcal{A}(v_3)| \geq 4$. There exists no $(1, \alpha, w, w_0)$-critical path for every $\alpha \in \mathcal{A}(w_1)\cup\mathcal{A}(v_3)$, thus there exists a $(3, \alpha, w, w_0)$-critical path, and then $\mathcal{A}(w_1)\cup\mathcal{A}(v_3) \subseteq \Upsilon(ww_3)$. Hence, $\deg_G(w_3) \geq |\mathcal{A}(w_1)|+|\mathcal{A}(v_3)|+|\{3, \kappa-\Delta+1\}| \geq 6$.

Suppose that $4 \notin \Upsilon(w_0w_2)$ and there exists no $(\kappa-\Delta+1, 4, w_0, w_2)$-critical path. Reassigning 4 to w_0w_2 results in a new acyclic edge coloring ϱ_3 of $G - ww_0$. Similarly, we can prove $\deg_G(w_4) \geq 6$ by replacing ϕ with ϱ_3.

If $4 \in \Upsilon(w_0w_2)$, then reassigning 1 to w_0w_2 and reassigning 4 to w_0v_3 will take us back to Subcase 2.1.3. If there exists a $(\kappa-\Delta+1, 4, w_0, w_2)$-critical path and $4 \in \Upsilon(w_0w_1)$, then reassigning 1 to w_0w_2 and 4 to w_0v_3 will take us back to Subcase 2.1.3 again.

Hence, we have that $\deg_G(w_4) \geq 6$. By symmetry, we also have that $\deg_G(w_i) \geq 6$ for $4 \leq i \leq \deg_G(w) - 1$.

Subcase 2.1.5. $\{\lambda_1, \lambda_2\} \cap \{1, 2\} = \emptyset$ and the color on w_0v_3 is a common color.

Without loss of generality, assume that $\phi(w_0w_1) = \kappa-\Delta+1$, $\phi(w_0w_2) = 3$, $\phi(w_0v_3) = 4$. If $1 \notin \Upsilon(w_0w_2) \cup \Upsilon(w_0v_3)$, then reassigning 1 to w_0w_2 will take us back to Subcase 2.1.3. Hence, $1 \in \Upsilon(w_0w_2) \cup \Upsilon(w_0v_3)$ and 1 appears at least twice in $\mathbb{S}$. If $2 \notin \Upsilon(w_0w_1) \cup \Upsilon(w_0v_3)$, then reassigning 2 to w_0w_1 and β_1 to w_0w_2 will take us back to Subcase 2.1.3. Therefore, we have

$$\sum_{\alpha\in\mathcal{U}(w)\cup\mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) + |X| \geq |\mathcal{U}(w)| + 2 + |X| + |\{1, 2\}| \geq \deg_G(w) + 3.$$

Subcase 2.1.6. $\{\lambda_1, \lambda_2\} \cap \{1, 2\} = \emptyset$ and the color on w_0v_3 is not a common color.

Without loss of generality, assume that $\phi(w_0w_1) = 3$, $\phi(w_0w_2) = 4$, $\phi(w_0v_3) = \kappa-\Delta+1$.

Suppose that $1 \notin \Upsilon(w_0w_2) \cup \Upsilon(w_0v_3)$. Thus, there exists a $(3, 1, w_0, w_2)$-critical path; otherwise, reassigning 1 to w_0w_2 and α_1 to ww_0 results in an acyclic edge coloring of G. But reassigning α_1, β_1 and 1 to ww_0, w_0w_2 and w_0v_3 respectively, yields an acyclic edge coloring of G. Hence, $1 \in \Upsilon(w_0w_2) \cup \Upsilon(w_0v_3)$ and 1 appears at least twice in $\mathbb{S}$. Similarly, we have that $2 \in \Upsilon(w_0w_1) \cup \Upsilon(w_0v_3)$. Therefore, we have

$$\sum_{\alpha\in\mathcal{U}(w)\cup\mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\alpha) + |X| \geq |\mathcal{U}(w)| + 2 + |X| + |\{1, 2\}| \geq \deg_G(w) + 3.$$

Subcase 2.2. $|\mathcal{U}(w) \cap \mathcal{U}(w_0)| = 3$.

Claim 6. Every color in $\mathcal{U}(w)$ is in $\mathbb{S}$.

Proof. Assume that w_0v_1, w_0v_2 and w_0v_3 are colored with λ_1, λ_2 and λ_3, respectively. Suppose that $\lambda_1 \notin \mathbb{S}$. If there is no $(\lambda_2, \alpha_1, w_0, v_1)$-critical path, then reassigning α_1 and λ_1 to w_0v_1 and w_0v_3 respectively, results in a new acyclic edge coloring of $G - ww_0$, which contradicts $(*)$. Hence, there exists a $(\lambda_2, \alpha_1, w_0, v_1)$-critical path, and hence there exists a $(\lambda_3, \alpha_1, w_0, w)$-critical path. But reassigning α_1 and λ_1 to w_0v_1 and w_0v_2, yields another acyclic edge coloring of $G - ww_0$, which contradicts $(*)$.

Hence, we have that $\lambda_1 \in \mathbb{S}$. By symmetry, we have that $\{\lambda_1, \lambda_2, \lambda_3\} \subseteq \mathbb{S}$. Let τ be an arbitrary color in $\mathcal{U}(w) \setminus (\mathbb{S} \cup \{\lambda_1, \lambda_2, \lambda_3\})$. Let σ be obtained from ϕ by reassigning τ to w_0v_1. It is obvious that σ is an acyclic edge coloring of $G - ww_0$. So we can obtain a similar contradiction by replacing ϕ with σ. So we conclude that $\mathcal{U}(w) \subseteq \mathbb{S}$. □

$$\sum_{\theta\in\mathcal{U}(w)\cup\mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq |\mathcal{U}(w)| = \deg_G(w) - 1,$$

In the following discussion, suppose that $\deg_G(w) + \deg_G(w_0) \leq \kappa - \Delta + 5$ and ww_0 is contained in two triangles ww_0w_1 and ww_0w_2 ($w_1 = v_1$ and $w_2 = v_2$).

Subcase 2.2.1. $\mathcal{U}(w_0) \cap \{1,2\} = \{1,2\}$.

By symmetry, assume that $\phi(w_0w_1) = 3, \phi(w_0w_2) = 1, \phi(w_0v_3) = 2$. Since $\alpha_1 \notin \mathcal{U}(w_1)$, it follows that there exists a $(2, \alpha_1, w_0, w)$-critical path. Reassigning α_1 to w_0w_1 will take us back to Subcase 2.1.2.

Subcase 2.2.2. $\mathcal{U}(w_0) \cap \{1,2\} = \{\lambda^*\}$ and λ^* is not on w_0v_3.

By symmetry, assume that $\phi(w_0w_1) = 3, \phi(w_0w_2) = 1, \phi(w_0v_3) = 5$. Since $\alpha_1 \notin \mathcal{U}(w_1)$, it follows that there exists a $(5, \alpha_1, w_0, w)$-critical path. Reassigning α_1 to w_0w_1 will take us back to Subcase 2.1.3.

Subcase 2.2.3. $\mathcal{U}(w_0) \cap \{1,2\} = \{\lambda^*\}$ and λ^* is on w_0v_3.

By symmetry, assume that $\phi(w_0w_1) = 3, \phi(w_0w_2) = 4, \phi(w_0v_3) = 1$. Since $\alpha_1 \notin \mathcal{U}(w_1)$, it follows that there exists a $(4, \alpha_1, w_0, w)$-critical path. Reassigning α_1 to w_0w_1 will take us back to Subcase 2.1.4.

Subcase 2.2.4. $\mathcal{U}(w_0) \cap \{1,2\} = \emptyset$.

By symmetry, assume that $\phi(w_0w_1) = 3, \phi(w_0w_2) = 4, \phi(w_0v_3) = 5$. Suppose that 1 only appears once in $\mathbb{S}$. Reassigning 1 to w_0w_2 will create a (3, 1)-dichromatic cycle containing w_0w_2, for otherwise, we go back to Subcase 2.2.2. But Reassigning 1 to w_0v_3 will take us back to Subcase 2.2.3. Hence, the color 1 appears at least twice in $\mathbb{S}$. Similarly, the color 2 appears at least twice in $\mathbb{S}$. Hence, we have

$$\sum_{\theta\in\mathcal{U}(w)\cup\mathcal{U}(w_0)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq \deg_G(w) - 1 + |\{1,2\}| = \deg_G(w) + 1. \qquad \square$$

3.3 Local structure on 5-vertices

Lemma 10. Let G be a κ-deletion-minimal graph with $\kappa \geq \Delta + 5$ and let u be a 5-vertex.

(a) If u is contained in a triangle wuw_1w with $\deg_G(w) \leq \kappa - \Delta$ and $\deg_G(w_1) \leq 6$, then

$$\sum_{x\in N_G(u)} \deg_G(x) \geq 2\kappa - \deg_G(u) + 12 = 2\kappa + 7. \tag{5}$$

(b) If u is contained in a triangle wuw_1w with $\deg_G(w) \leq \kappa - \Delta - 1$ and $\deg_G(w_1) \leq 7$, then

$$\sum_{x\in N_G(u)} \deg_G(x) \geq 2\kappa - \deg_G(u) + 12 = 2\kappa + 7. \tag{6}$$

Proof. We may assume that

(*) The graph $G - wu$ admits an acyclic edge coloring ϕ such that the number of common colors at w and u is minimum.

Here, (a) and (b) will be proved together, so we may assume that $\deg_G(w) \leq \kappa - \Delta$. Since $\deg_G(w) + \deg_G(u) \leq \kappa - \Delta + 5 < \kappa + 2$, we have that $|\Upsilon(wu) \cap \Upsilon(uw)| = m \geq 1$. It follows that $|C(wu)| = \kappa - (\deg_G(w) + \deg_G(u) - m - 2) \geq \Delta - 2$. Without loss of generality, let $N_G(w) = \{u, w_1, w_2, \dots\}$ and $\phi(ww_i) = i$ for $1 \leq i \leq \deg_G(w) - 1$. Let $N_G(u) = \{w, u_1, u_2, u_3, u_4\}$ and $\mathbb{S} = \Upsilon(uu_1) \uplus \Upsilon(uu_2) \uplus \Upsilon(uu_3) \uplus \Upsilon(uu_4)$.

Let $\mathcal{A}(u_1) = C(wu) \setminus \mathcal{U}(u_1) = \{\alpha_1, \alpha_2, \dots\}$, $\mathcal{A}(u_2) = C(wu) \setminus \mathcal{U}(u_2) = \{\beta_1, \beta_2, \dots\}$, $\mathcal{A}(u_3) = C(wu) \setminus \mathcal{U}(u_3) = \{\xi_1, \xi_2, \dots\}$, $\mathcal{A}(u_4) = C(wu) \setminus \mathcal{U}(u_4) = \{\zeta_1, \zeta_2, \dots\}$ and $\mathcal{A}(w_2) = C(wu) \setminus \mathcal{U}(w_2) = \{\zeta_1^*, \zeta_2^*, \dots\}$.

Claim 1. For every color θ in $C(wu)$, there exists an (λ, θ, u, w)-critical path for some $\lambda \in \mathcal{U}(w) \cap \mathcal{U}(u)$. Consequently, $\mathrm{mul}_{\mathbb{S}}(\theta) \geq 1$.

Case 1. $\mathcal{U}(w) \cap \mathcal{U}(u) = \{\lambda\}$. By symmetry, we may assume that $w_1 = u_1$.

It follows that $|C(wu)| = \kappa - (\deg_G(w) + \deg_G(u) - 3) \geq \Delta - 2$.

Subcase 1.1. The edge ww_1 is colored with λ. By symmetry, assume that $\phi(uw_1) = \kappa-\Delta+2$, $\phi(uu_2) = 1$, $\phi(uu_3) = \kappa-\Delta$, $\phi(uu_4) = \kappa - \Delta + 1$.

By Claim 1, we have that $\{\kappa-\Delta+3, \ldots, \kappa\} \subseteq \Upsilon(ww_1) \cap \Upsilon(uu_2)$. Moreover, $\mathcal{U}(w_1) = \{1, \kappa-\Delta+2\} \cup \{\kappa-\Delta+3, \ldots, \kappa\}$, $\deg_G(w_1) = \Delta$ and $\deg_G(u_2) \geq \Delta - 1$. Notice that $|\Upsilon(uu_2) \cap \{2, 3, \ldots, \kappa - \Delta - 1\}| \leq 1$, thus there exists a color ζ which is in $\{2, 3, \ldots, \kappa - \Delta - 1\} \setminus \Upsilon(uu_2)$ (note that this set is nonempty). But assigning $\kappa - \Delta + 2$ to uw and ζ to uw_1 results in an acyclic edge coloring of G, a contradiction.

Subcase 1.2. The edge uw_1 is colored with λ. By symmetry, assume that $\phi(uw_1) = 2$, $\phi(uu_2) = \kappa-\Delta$, $\phi(uu_3) = \kappa-\Delta+1$, $\phi(uu_4) = \kappa - \Delta + 2$.

By Claim 1, we have that $\{\kappa - \Delta + 3, \ldots, \kappa\} \subseteq \Upsilon(uw_1) \cap \Upsilon(ww_2)$ and $\deg_G(w_1) = \Delta$ and $\deg_G(w_2) \geq \Delta - 1$. Modify ϕ by reassigning 1 to wu and reassigning a color in $\{\kappa - \Delta, \kappa - \Delta + 1, \kappa - \Delta + 2\} \setminus \mathcal{U}(w_2)$ to ww_1, we obtain an acyclic edge coloring of G, a contradiction.

Subcase 1.3. Neither w_1w nor w_1u is colored with λ. By symmetry, assume that $\phi(uw_1) = \kappa - \Delta$, $\phi(uu_2) = 2$, $\phi(uu_3) = \kappa - \Delta + 1$, $\phi(uu_4) = \kappa - \Delta + 2$.

By Claim 1, we have that $C(wu) \subseteq \Upsilon(uu_2) \cap \Upsilon(ww_2)$ and $\deg_G(w_2) \geq \Delta - 1$ and $\deg_G(u_2) \geq \Delta - 1$. Notice that $\{1, \kappa - \Delta\} \nsubseteq \mathcal{U}(w_2)$.

If $\deg_G(w) \leq \kappa - \Delta - 1$, then $\mathcal{U}(w) = \{1, 2, \ldots, \kappa - \Delta - 2\}$ and $\deg(w_2) = \deg(u_2) = \Delta$, but reassigning 1 to uu_2 will take us back to Subcase 1.1.

So we may assume that $\deg(w) = \kappa - \Delta$, $C(wu) = \{\kappa - \Delta + 3, \ldots, \kappa\}$ and $\deg(w_1) \leq 6$.

Suppose that $C(wu) \subseteq \mathcal{U}(w_1)$. Thus $\mathcal{U}(w_1) = \{1, \kappa - \Delta\} \cup C(wu)$. If $1 \notin \mathcal{U}(w_2)$, then reassigning $1, \kappa - \Delta$ and 3 to wu, ww_1 and w_1u respectively results in an acyclic edge coloring of G, a contradiction. So we may assume that $1 \in \mathcal{U}(w_2)$ and $\Upsilon(ww_2) = \{1\} \cup \{\kappa - \Delta + 3, \ldots, \kappa\}$. But reassigning $\kappa - \Delta$ to wu and 3 to w_1u results in an acyclic edge coloring of G, a contradiction. Hence, we have that $C(wu) \nsubseteq \mathcal{U}(w_1)$.

We further suppose that $1 \in \mathcal{U}(u_2)$ and $\Upsilon(uu_2) = \{1\} \cup \{\kappa - \Delta + 3, \ldots, \kappa\}$. If there is a $(2, 1, u, w)$-critical path, then $\deg_G(w_2) = \Delta(G)$ and $\Upsilon(ww_2) = \{1\} \cup \{\kappa - \Delta + 3, \ldots, \kappa\}$, but reassigning $\kappa - \Delta$ to ww_2 will take us back to Subcase 1.2. So we may assume that there is no $(2, 1, u, w)$-critical path. There exists a $(\tau^*, \alpha_1, w, w_1)$-critical path with some $\tau^* \in \mathcal{U}(w) \setminus \{1, 2\}$, otherwise reassigning α_1 to ww_1 and 1 to uw will result in an acyclic edge coloring of G. By symmetry, assume that $\tau^* = 3$ and there exists a $(3, \alpha_1, w, w_1)$-critical path. But reassigning 3 to uu_2 and α_1 to wu results in an acyclic edge coloring of G.

So we may assume that $1 \notin \mathcal{U}(u_2)$. There exists a $(\kappa - \Delta + 1, 1, u, u_2)$- or $(\kappa - \Delta + 2, 1, u, u_2)$-critical path; otherwise, reassigning 1 to uu_2 will take us back to Subcase 1.1. By symmetry, assume that there exists a $(\kappa-\Delta+2, 1, u, u_2)$-critical path and $1 \in \Upsilon(uu_4)$, thus $\deg_G(u_2) = \Delta(G)$ and $\Upsilon(uu_2) = \{\kappa - \Delta + 2, \kappa - \Delta + 3, \ldots, \kappa\}$.

There exists a $(\kappa - \Delta + 1, \alpha_1, u, w_1)$- or $(\kappa - \Delta + 2, \alpha_1, u, w_1)$-critical path; otherwise, reassigning α_1 to uw_1 and $\kappa - \Delta$ to uw will result in an acyclic edge coloring of G. Hence, $\{\kappa - \Delta + 1, \kappa - \Delta + 2\} \cap \mathcal{U}(w_1) \neq \emptyset$. Similarly, there exists a (τ, α_1, w, w_1)-critical path with some $\tau \in \mathcal{U}(w) \setminus \{1, 2\}$. By symmetry, assume that $\tau = 3$ and there exists a $(3, \alpha_1, w, w_1)$-critical path. Hence, $|\mathcal{U}(w_1) \cap (\mathcal{U}(w) \cup \mathcal{U}(u))| \geq 4$ and $|\mathcal{U}(w_1) \cap C(wu)| \leq 2$, and then $|C(wu) \setminus \mathcal{U}(w_1)| \geq \Delta - 4$.

Suppose that $\mathcal{A}(u_4) \cap \mathcal{A}(w_1) \neq \emptyset$, say $\zeta \in \mathcal{A}(u_4) \cap \mathcal{A}(w_1)$. Thus there exists a $(\kappa - \Delta + 1, \zeta, u, w_1)$-critical path; otherwise, reassigning ζ to uw_1 and $\kappa - \Delta$ to uw will result in an acyclic edge coloring of G. There exists a $(2, \kappa - \Delta + 2, u, w)$-critical path, otherwise reassigning ζ to uu_4 and $\kappa - \Delta + 2$ to uw will result in an acyclic edge coloring of G. Hence, we have that $\Upsilon(ww_2) = \Upsilon(uu_2) = \{\kappa - \Delta + 2, \kappa - \Delta + 3, \ldots, \kappa\}$. But reassigning $\kappa - \Delta$ to ww_2 will take us back to Subcase 1.2. So we have that $\mathcal{A}(u_4) \cap \mathcal{A}(w_1) = \emptyset$.

There exists a $(\kappa - \Delta + 2, 3, u, u_2)$-critical path, for otherwise reassigning 3 to uu_2 and α_1 to uw will result in an acyclic edge coloring of G. It follows that $\{1, 3\} \cup \mathcal{A}(w_1) \subseteq \Upsilon(uu_4)$. If $2 \notin \mathcal{U}(u_4)$ and there exists no $(\kappa-\Delta+1, 2, u, u_4)$-critical path, then reassigning 2, 1 and α_1 to uu_4, uu_2 and uw, respectively, will result in an acyclic edge coloring of G. Hence, $\mathcal{U}(u_4) = \{1, 2, 3, \kappa - \Delta + 2\} \cup \mathcal{A}(w_1)$ or $\mathcal{U}(u_4) = \{1, 3, \kappa - \Delta + 1, \kappa - \Delta + 2\} \cup \mathcal{A}(w_1)$. Consequently, we have that $|\mathcal{A}(w_1)| = \Delta - 4$, and then $\mathcal{U}(w_1) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, 3, \kappa - \Delta, \kappa - \Delta + 1\}$ or $\{1, 3, \kappa - \Delta, \kappa - \Delta + 2\}$.

There exists a $(\kappa - \Delta + 1, 2, u, w_1)$- or $(\kappa - \Delta + 2, 2, u, w_1)$-critical path, for otherwise we reassign $\alpha_1, 2$ and $\kappa - \Delta$ to uw, uw_1 and uu_2. Thus, $\mathcal{U}(u_4) = \{1, 2, 3, \kappa - \Delta + 2\} \cup \mathcal{A}(w_1)$. If $\mathcal{U}(w_1) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, 3, \kappa - \Delta, \kappa - \Delta + 2\}$, then we reassign $\alpha_1, 2, 3$ and $\kappa - \Delta$ to uw, uw_1, uu_2 and uu_4. Therefore, $\mathcal{U}(w_1) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, 3, \kappa - \Delta, \kappa - \Delta + 1\}$, but reassigning $\kappa - \Delta + 2, 1$ and 4 to uw, uu_2 and uu_4 results in an acyclic edge coloring of G.

Case 2. $\mathcal{U}(w) \cap \mathcal{U}(u) = \{\lambda_1, \ldots, \lambda_m\}$ and $m \geq 2$.

We can relabel the vertices in $\{u_1, u_2, u_3, u_4\}$ as $\{v_1, v_2, v_3, v_4\}$. By symmetry, we may assume that $\phi(uv_i) = \lambda_i$ for $i \in \{1, \ldots, m\}$.

Claim 2. The sets $\mathcal{A}(v_1), \mathcal{A}(v_2), \ldots, \mathcal{A}(v_m)$ are pairwise disjoint.

Proof. Suppose, to the contrary, that $\alpha \in \mathcal{A}(v_1) \cap \mathcal{A}(v_2)$. By Claim 1 and the symmetry, we may assume that there exists a $(\lambda_3, \alpha, u, w)$-critical path and $m \geq 3$, which implies that there exists no $(\lambda_3, \alpha, u, v_2)$-critical path. Consequently, there exists a $(\phi(uv_4), \alpha, u, v_2)$-critical path; otherwise, reassigning α to uv_2 to obtain a new acyclic edge coloring of $G - wu$, which contradicts the minimality of m. Now, reassigning α to uv_1 to obtain an acyclic edge coloring π of $G - wu$, but $|\mathcal{U}_\pi(u) \cap \mathcal{U}_\pi(w)| < |\mathcal{U}(u) \cap \mathcal{U}(w)|$, which is a contradiction. □

Claim 3. Every color in $C(wu)$ appears at least twice in $\mathbb{S}$.

Proof. Suppose that there exists a color α in $C(wu)$ such that $\mathrm{mul}_{\mathbb{S}}(\alpha) = 1$. By Claim 1 and symmetry, we may assume that there exists a $(\lambda_1, \alpha, u, w)$-critical path and $\alpha \in \mathcal{U}(v_1)$. But reassigning α to uv_2 results in a new acyclic edge coloring of $G - wu$, which contradicts the assumption $(*)$. □

Let $X = \{\, \theta \mid \theta \in C(wu) \text{ and } \mathrm{mul}_{\mathbb{S}}(\theta) \geq 3 \,\}$.

$$
\begin{aligned}
&\sum_{x \in N_G(u)} \deg_G(x) \\
&= \deg_G(u) + \deg_G(w) - 1 + \sum_{\theta \in [\kappa]} \mathrm{mul}_{\mathbb{S}}(\theta) \\
&= \deg_G(u) + \deg_G(w) - 1 + \sum_{\theta \in C(wu)} \mathrm{mul}_{\mathbb{S}}(\theta) + \sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) \\
&\geq \deg_G(u) + \deg_G(w) - 1 + 2|C(wu)| + |X| + \sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) \\
&= \deg_G(u) + \deg_G(w) - 1 + 2(\kappa - (\deg_G(w) + \deg_G(u) - 2 - m)) + |X| + \sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) \\
&= 2\kappa - \deg_G(u) - \deg_G(w) + 2m + 3 + |X| + \sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta)
\end{aligned}
$$

It is sufficient to prove that

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq \deg_G(w) - 2m + 9. \tag{7}$$

Subcase 2.1. $\mathcal{U}(w) \cap \mathcal{U}(u) = \{\lambda_1, \lambda_2\}$ and $w_1 = u_1$. Note that $\mathcal{A}(w_1) \neq \emptyset$.

Subcase 2.1.1. The two colors on the edges w_1w and w_1u are all common colors.

Without loss of generality, assume that $\phi(uw_1) = 2$, $\phi(uu_2) = 1$, $\phi(uu_3) = \kappa - \Delta$ and $\phi(uu_4) = \kappa - \Delta + 1$. Consequently, we conclude that $\{\kappa - \Delta + 2, \ldots, \kappa\} \subseteq \mathcal{U}(w_1)$ and $\deg_G(w_1) \geq \Delta + 1$, a contradiction.

Subcase 2.1.2. The color on w_1w is a common color and the color on w_1u is not a common color.

Without loss of generality, assume that $\phi(uw_1) = \kappa - \Delta$, $\phi(uu_2) = 1$, $\phi(uu_3) = 2$ and $\phi(uu_4) = \kappa - \Delta + 1$.

For every color $\alpha_i \in \mathcal{A}(w_1)$, there exists a $(\theta_i, \alpha_i, w, w_1)$-critical path with some $\theta_i \in \mathcal{U}(w) \setminus \{1, 2\}$; otherwise, reassigning α_i to ww_1 will take us back to Case 1. By symmetry, we may assume that there exists a $(3, \alpha^*, w, w_1)$-critical path with some α^*, and then $3 \in \mathcal{U}(w_1)$. If $\Upsilon(ww_2) \subseteq C(wu)$, then reassigning $\kappa - \Delta$ to ww_2 will take us back to Subcase 2.1.1. So we have that $\Upsilon(ww_2) \nsubseteq C(wu)$ and $\mathcal{A}(w_2) \neq \emptyset$. Consequently, for every color $\zeta_i^* \in \mathcal{A}(w_2)$, there exists a $(\mu_i, \zeta_i^*, w, w_2)$-critical path with some $\mu_i \in \mathcal{U}(w) \setminus \{1, 2\}$; otherwise, reassigning ζ_i^* to ww_2 will take us back to Case 1. Hence, $\{1, 3, \kappa - \Delta\} \subseteq \mathcal{U}(w_1)$ and $\{2, \mu_1\} \subseteq \mathcal{U}(w_2)$, and then $|\mathcal{A}(w_1)| \geq 2$ and $|\mathcal{A}(w_2)| \geq 1$.

If $\Upsilon(uu_2) \subseteq C(wu)$, then reassigning μ_1 to uu_2 and ζ_1^* to wu results in an acyclic edge coloring of G, a contradiction. Thus, we have that $\Upsilon(uu_2) \nsubseteq C(wu)$ and $\mathcal{A}(u_2) \neq \emptyset$. For every color $\beta_i \in \mathcal{A}(u_2)$, there exists an $(\varepsilon_i, \beta_i, u, u_2)$-critical path with some $\varepsilon_i \in \{\kappa - \Delta, \kappa - \Delta + 1\}$; otherwise, reassigning β_i to uu_2 will take us back to Case 1.

If $\Upsilon(uu_3) \subseteq C(wu)$, then reassigning 3 to uu_3 and α^* to wu results in an acyclic edge coloring of G, a contradiction. Thus, we have that $\Upsilon(uu_3) \nsubseteq C(wu)$ and $\mathcal{A}(u_3) \neq \emptyset$. Consequently, for every color $\xi_i \in \mathcal{A}(u_3)$, there exists a (m_i, ξ_i, u, u_3)-critical path with some $m_i \in \{\kappa - \Delta, \kappa - \Delta + 1\}$; otherwise, reassigning ξ_i to uu_3 will take us back to Case 1.

Claim 4. $\mathcal{A}(u_2) \cap \mathcal{A}(u_4) = \emptyset$.

Proof of Claim 4. Suppose that $\beta_1 \in \mathcal{A}(u_2) \cap \mathcal{A}(u_4)$. It follows that there exists a $(\kappa - \Delta, \beta_1, u, u_2)$-critical path, $\kappa - \Delta \in \Upsilon(uu_2)$ and $\beta_1 \in \Upsilon(uw_1)$. Also, there exists a $(1, \kappa - \Delta + 1, w, u)$- or $(2, \kappa - \Delta + 1, w, u)$-critical path; otherwise, reassigning β_1 to uu_4 and $\kappa - \Delta + 1$ to wu results in an acyclic edge coloring of G. Suppose that there exists a $(1, \kappa - \Delta + 1, w, u)$-critical path and $\kappa - \Delta + 1 \in \Upsilon(ww_1) \cap \Upsilon(uu_2)$. It follows that $\{1, \kappa - \Delta, \kappa - \Delta + 1\} \subseteq \mathcal{U}(u_2)$ and $|\mathcal{A}(u_2)| \geq 2$. Furthermore, we can conclude that $\{1, 3, \kappa-\Delta, \kappa-\Delta+1, \beta_1\} \cup \mathcal{A}(w_2) \subseteq \mathcal{U}(w_1)$. Note that $\mathcal{A}(w_2) \cap \mathcal{A}(u_2) = \emptyset$, thus $\mathcal{U}(w_1) = \{1, 3, \kappa - \Delta, \kappa - \Delta + 1, \beta_1\} \cup \mathcal{A}(w_2)$ and $\mathcal{U}(w_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{2, \mu_1\}$. Recall that $\beta_2 \notin \mathcal{A}(w_2)$, thus $\varepsilon_2 = \kappa - \Delta + 1$ and there exists a $(\kappa - \Delta + 1, \beta_2, u, u_2)$-critical path. Now, reassigning β_2 to uw_1 and $\kappa - \Delta$ to wu results in an acyclic edge coloring of G.

So, we may assume that there exists a $(2, \kappa - \Delta + 1, w, u)$-critical path. Hence, $\{1, \kappa - \Delta, 3, \beta_1\} \cup \mathcal{A}(w_2) \subseteq \mathcal{U}(w_1)$ and $\{2, \mu_1, \kappa-\Delta+1\} \subseteq \Upsilon(ww_2)$. It follows that $\mathcal{U}(w_1) = \{1, \kappa-\Delta, 3, \beta_1\} \cup \mathcal{A}(w_2)$ and $\mathcal{U}(w_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{2, \mu_1, \kappa-\Delta+1\}$. Now, reassigning α_1 to uw_1 and $\kappa - \Delta$ to wu results in an acyclic edge coloring of G. This completes the proof of Claim 4. □

Claim 5. $\mathcal{A}(u_2) \cap \mathcal{A}(w_1) = \emptyset$.

Proof of Claim 5. By contradiction, assume that $\alpha_1 = \beta_1$. It follows that there exists a $(\kappa - \Delta + 1, \beta_1, u, u_2)$-critical path and $\kappa - \Delta + 1 \in \mathcal{U}(u_2)$. There exists a $(2, \kappa - \Delta, w, u)$-critical path; otherwise, reassigning β_1 to uw_1 and $\kappa - \Delta$ to uw results in an acyclic edge coloring of G, a contradiction. So we have that $\kappa - \Delta \in \Upsilon(ww_2) \cap \Upsilon(uu_3)$.

Note that $\{1, \kappa-\Delta, 3\} \subseteq \mathcal{U}(w_1)$ and $\{2, \mu_1, \kappa-\Delta\} \subseteq \mathcal{U}(w_2)$. If $\deg_G(w) \leq \kappa-\Delta$ and $\deg_G(w_1) \leq 6$, then $\mathcal{A}(w_2) \subseteq \mathcal{U}(w_1)$ with $|\mathcal{A}(w_2)| \geq 2$, and then $|\mathcal{U}(w_1) \cap \mathcal{U}(w)| \leq 3$; similarly, if $\deg_G(w) \leq \kappa-\Delta-1$ and $\deg_G(w_1) \leq 7$, then $\mathcal{A}(w_2) \subseteq \mathcal{U}(w_1)$ with $|\mathcal{A}(w_2)| \geq 3$, and then $|\mathcal{U}(w_1) \cap \mathcal{U}(w)| \leq 3$. If $\mathcal{U}(w_1) \cap \mathcal{U}(w) = \{1, 3\}$, then $\{2, \kappa - \Delta\} \subseteq \mathcal{U}(u_3) \cap (\mathcal{U}(w) \cup \mathcal{U}(u))$; otherwise, reassigning α_1, α_2 and 3 to uw_1, uw and uu_3 respectively results in an acyclic edge coloring of G. Suppose that $\mathcal{U}(w_1) \cap \mathcal{U}(w) = \{1, 3, s\}$. Since $|\mathcal{A}(w_1)| \geq 3$, thus there exists a $\tau \in \{3, s\}$ and α_i, α_j such that both (τ, α_i, w, w_1)- and (τ, α_j, w, w_1)-critical path exist, and thus $\{2, \kappa - \Delta\} \subseteq \mathcal{U}(u_3) \cap (\mathcal{U}(w) \cup \mathcal{U}(u))$; otherwise, reassigning α_i, α_j and τ to uw_1, uw and uu_3 respectively. Anyway, we have that $|\mathcal{U}(u_3) \cap (\mathcal{U}(w) \cup \mathcal{U}(u))| \geq 3$.

If $\kappa - \Delta$ only appears only once (at u_3) in $\mathbb{S}$, then reassigning $\kappa - \Delta$ to uu_2 and β_1 to uw_1 will take us back to Case 1. So we conclude that the color $\kappa - \Delta$ appears at least twice in $\mathbb{S}$.

If $1 \notin \mathbb{S} \setminus \mathcal{U}(w_1)$, then reassigning $1, \beta_1$ and ξ_1 to uu_4, uu_2 and wu respectively, results in an acyclic edge coloring of G, a contradiction. Therefore, the color 1 appears at least twice in $\mathbb{S}$.

Suppose that $4 \notin \mathbb{S}$. Thus there exists a $(4, \xi_1, w, u_2)$-alternating path; otherwise, reassigning 4 to uu_2 and ξ_1 to wu results in an acyclic edge coloring of G, a contradiction. Now, reassigning $4, \beta_1$ and ξ_1 to uu_4, uu_2 and wu respectively, results in an acyclic edge coloring G, a contradiction. So we conclude that $4 \in \mathbb{S}$. By symmetry, we can also obtain that every color in $\mathcal{U}(w) \setminus \{1, 2, 3\}$ appears in $\mathbb{S}$.

Suppose that every color in $\mathcal{U}(w) \setminus \{1, 2\}$ appears exactly once in $\mathbb{S}$. Suppose that $\mathcal{U}(w_1) \cap (\mathcal{U}(w) \setminus \{1, 2\}) = \{3, s\}$. Thus, $\mathcal{U}(w_1) = \{1, \kappa - \Delta, 3, s\} \cup \mathcal{A}(w_2)$ and $\kappa - \Delta + 1 \notin \mathcal{U}(w_1)$. Since $|\mathcal{A}(w_1)| \geq 3$, thus there exists a $\tau \in \{3, s\}$ and α_i, α_j such that both (τ, α_i, w, w_1)- and (τ, α_j, w, w_1)-critical path exist. Reassigning τ, α_i and α_j to uu_3, uw_1 and wu respectively, results in an acyclic edge coloring of G, a contradiction. So we may assume that $|\mathcal{U}(w_1) \cap (\mathcal{U}(w) \setminus \{1, 2\})| = 1$, that is $\mathcal{U}(w_1) \cap (\mathcal{U}(w) \setminus \{1, 2\}) = \{3\}$. Reassigning $3, \alpha_1$ and α_2 to uu_3, uw_1 and wu respectively, results in an acyclic edge coloring of G, a contradiction. Hence, we may assume that the color 3 appears at least twice in $\mathbb{S}$.

Suppose that $\xi_1 \in \mathcal{A}(u_4)$. Thus, there exists a $(\kappa - \Delta, \xi_1, u, u_3)$-critical path; otherwise, reassigning ξ_1 to uu_3 will take us back to Case 1. Furthermore, $\kappa - \Delta + 1 \in \Upsilon(ww_1) \cup \Upsilon(uu_3)$; otherwise, reassigning ξ_1 to uu_4 and $\kappa - \Delta + 1$ to wu results in an acyclic edge coloring of G. If $2 \notin \mathbb{S}$, then reassigning $\alpha_1, 2$ and ξ_1 to wu, uw_1 uu_3 respectively, results in an acyclic edge coloring of G. So we have that $2 \in \mathbb{S}$. Hence,

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq |\{4, \ldots, \deg_G(w) - 1\}| + 2|\{1, 3, \kappa - \Delta, \kappa - \Delta + 1\}| + |\{2\}| = \deg_G(w) + 5.$$

So we may assume that $\mathcal{A}(u_3) \cap \mathcal{A}(u_4) = \emptyset$. It is obvious that $\mathcal{A}(u_3) \subseteq X$. Hence,

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq |\{4, \ldots, \deg_G(w) - 1\}| + 2|\{1, 3, \kappa - \Delta\}| + |\{\kappa - \Delta + 1\}| + |\mathcal{A}(u_3)| \geq \deg_G(w) + 4.$$

The equality holds only if $\kappa - \Delta + 1$ appears only once in $\mathbb{S}$ and 2 does not appear in $\mathbb{S}$; but reassigning $\alpha_1, 2$ and ξ_1 to wu, uw_1 and uu_3 respectively, results in an acyclic edge coloring. Therefore, inequality (7) holds, we are done. This completes the proof Claim 5. □

By Claim 5, the three sets $\mathcal{A}(w_1), \mathcal{A}(u_2)$ and $\mathcal{A}(u_3)$ are pairwise disjoint.

(1) Suppose that there exists no $(2, \kappa-\Delta, w, u)$-critical path. This implies that there exists a $(\kappa-\Delta+1, \alpha_i, u, w_1)$-critical path; otherwise, reassigning α_i to uw_1 and $\kappa-\Delta$ to wu results in an acyclic edge coloring of G. Thus, $\{1, 3, \kappa-\Delta, \kappa-\Delta+1\} \subseteq \mathcal{U}(w_1)$ and $\mathcal{A}(w_1) \subseteq \mathcal{U}(u_4)$. Note that $\mathcal{A}(u_2) \cup \mathcal{A}(u_3) \subseteq \mathcal{U}(w_1)$, thus $|\mathcal{A}(u_2)| = |\mathcal{A}(u_3)| = 1$ and $\mathcal{U}(w_1) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, 3, \kappa-\Delta, \kappa-\Delta+1\}$. Similarly, we know that $\mathcal{A}(u_2) \cup \mathcal{A}(w_2) \subseteq \mathcal{U}(w_1)$, which implies that $|\mathcal{A}(w_2)| = |\mathcal{A}(u_3)| = 1$ and $\mathcal{A}(w_2) = \mathcal{A}(u_3)$. Hence, $\mathcal{U}(w_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{2, \mu_1\}$. By Claim 4, we conclude that $\mathcal{U}(u_4) \supseteq \mathcal{A}(w_1) \cup \mathcal{A}(u_2) \cup \{\kappa-\Delta+1\}$. If $\Upsilon(uu_4) \subseteq C(uw)$, then reassigning $3, \alpha^*$ and $\kappa-\Delta$ to uu_4, uw_1 and wu respectively results in an acyclic edge coloring of G. Note that $|\mathcal{A}(w_1)| + |\mathcal{A}(u_2)| = \Delta - 2$, so we may assume that $|\Upsilon(uu_4) \cap (\mathcal{U}(w) \cup \mathcal{U}(u))| = 1$. In addition, $\Upsilon(uu_4) \cap C(uw) = \mathcal{A}(w_1) \cup \mathcal{A}(u_2)$ and $\mathcal{U}(u_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, \varepsilon_1\}$. Recall that $\mathcal{A}(w_1), \mathcal{A}(u_2)$ and $\mathcal{A}(u_3)$ are pairwise disjoint, thus $\mathcal{A}(u_3) \cap \mathcal{U}(u_4) = \emptyset$, and then there exists a $(\kappa-\Delta, \xi_1, u, u_3)$-critical path and $\kappa-\Delta \in \Upsilon(uu_3)$. There exists a $(1, \kappa-\Delta+1, u, w)$-critical path; otherwise, reassigning ξ_1 and $\kappa-\Delta+1$ to uu_4 and uw results in an acyclic edge coloring of G. Hence, $\mathcal{U}(u_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, \varepsilon_1\} = \{1, \kappa-\Delta+1\}$. There exists a $(\kappa-\Delta+1, \mu_1, u, u_2)$-critical path, otherwise, reassigning μ_1 to uu_2 and ζ_1^* to uw results in an acyclic edge coloring of G. Hence, $\Upsilon(uu_4) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{\mu_1, \kappa-\Delta+1\}$. Now, reassigning $\zeta_1^*, \alpha_1, \mu_1$ and $\kappa-\Delta$ to uw, uw_1, uu_2 and uu_4 respectively, yields an acyclic edge coloring of G.

(2) Now, we may assume that there exists a $(2, \kappa-\Delta, w, u)$-critical path and $\kappa-\Delta \in \Upsilon(uu_3) \cap \Upsilon(ww_2)$. Clearly, $\mathcal{U}(w_1) \supseteq \mathcal{A}(u_2) \cup \mathcal{A}(w_2) \cup \{1, 3, \kappa-\Delta\}$. If $\deg_G(w) \leq \kappa-\Delta-1$, then $\deg_G(w_1) \geq 2+3+3 = 8$, a contradiction. Thus, $\deg_G(w) = \kappa-\Delta$, which implies that $\mathcal{U}(w_1) = \mathcal{A}(u_2) \cup \mathcal{A}(w_2) \cup \{1, 3, \kappa-\Delta\}$, $|\mathcal{A}(u_2)| = 1$ and $|\mathcal{A}(w_2)| = 2$. It is easy to see that $\mathcal{U}(w_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{2, \kappa-\Delta, \mu_1\}$ and $\mathcal{U}(u_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, \varepsilon_1\}$. If $3 \notin \mathcal{U}(u_3)$, then there exists a $(\kappa-\Delta+1, 3, u, u_3)$-critical path; otherwise, reassigning α_1, α_2 and 3 to uw_1, uw and uu_3 respectively results in an acyclic edge coloring of G. Hence, we have that $\{3, \kappa-\Delta+1\} \cap \mathcal{U}(u_3) \neq \emptyset$ and $|\mathcal{A}(u_3)| \geq 2$. Recall that $\mathcal{A}(w_1), \mathcal{A}(u_2)$ and $\mathcal{A}(u_3)$ are disjoint, thus $\mathcal{U}(w_1) \supseteq \mathcal{A}(u_2) \cup \mathcal{A}(u_3) \cup \{1, 3, \kappa-\Delta\}$. Moreover, $\mathcal{U}(w_1) = \mathcal{A}(u_2) \cup \mathcal{A}(u_3) \cup \{1, 3, \kappa-\Delta\}$, $\mathcal{A}(u_3) = \mathcal{A}(w_2)$. If there exists a $\xi_i \notin \mathcal{U}(u_4)$, then there exists a $(\kappa-\Delta, \xi_i, u, u_3)$-critical path, and then reassigning ξ_i to uu_4 and $\kappa-\Delta+1$ to uw results in an acyclic edge coloring of G. So we have that $\mathcal{A}(u_2) \cup \mathcal{A}(u_3) \subseteq \mathcal{U}(u_4)$.

There exists a $(\kappa-\Delta, \mu_1, u, u_2)$- or $(\kappa-\Delta+1, \mu_1, u, u_2)$-critical path; otherwise, reassigning μ_1 to uu_2 and ζ_1^* to uw results in an acyclic edge coloring of G. If there exists a $(\kappa-\Delta, \mu_1, u, u_2)$-critical path, then $\mu_1 = 3$ and $\varepsilon_1 = \kappa-\Delta$; but reassigning μ_1, α^* and ζ_1^* to uu_2, uw_1 and uw results in an acyclic edge coloring. So there exists a $(\kappa-\Delta+1, \mu_1, u, u_2)$-critical path, thus $\varepsilon_1 = \kappa-\Delta+1$ and $\mathcal{U}(u_4) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{\mu_1, \kappa-\Delta+1\}$. Now, reassigning $\kappa-\Delta, \mu_1, \alpha^*$ and ζ_1^* to uu_4, uu_2, uw_1 and uw respectively, yields an acyclic edge coloring of G.

Subcase 2.1.3. The color on w_1w is not a common color and the color on w_1u is a common color.

Without loss of generality, assume that $\phi(uw_1) = 3$, $\phi(uu_2) = 2$, $\phi(uu_3) = \kappa-\Delta$ and $\phi(uu_4) = \kappa-\Delta+1$.

For every color $\alpha_i \in \mathcal{A}(w_1)$, there exists a $(\theta_i, \alpha_i, u, w_1)$-critical path with some $\theta_i \in \{\kappa-\Delta, \kappa-\Delta+1\}$; otherwise, reassigning α_i to uw_1 will take us back to Case 1. If $\Upsilon(uu_2) \subseteq C(wu)$, then reassigning 1 to uu_2 will take us back to Subcase 2.1.1. So we have that $\Upsilon(uu_2) \nsubseteq C(wu)$ and $\mathcal{A}(u_2) \neq \emptyset$. Consequently, for every color $\beta_i \in \mathcal{A}(u_2)$, there exists a $(\varepsilon_i, \beta_i, u, u_2)$-critical path with some $\varepsilon_i \in \{\kappa-\Delta, \kappa-\Delta+1\}$; otherwise, reassigning β_i to uu_2 will take us back to Case 1. Hence, we have $\{\kappa-\Delta, \kappa-\Delta+1\} \cap \Upsilon(uu_2) \neq \emptyset$.

Subcase 2.1.3.1. Suppose that $\{\kappa-\Delta, \kappa-\Delta+1\} \subseteq \mathcal{U}(w_1)$.

If $\{\kappa-\Delta, \kappa-\Delta+1\} \subseteq \mathcal{U}(u_2)$, then $\mathcal{U}(w_1) = \{1, 3, \kappa-\Delta, \kappa-\Delta+1\} \cup \mathcal{A}(u_2)$ and $\mathcal{U}(u_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{\kappa-\Delta, \kappa-\Delta+1, 2\}$; but reassigning α_1 to ww_1 and 1 to uw results in an acyclic edge coloring of G. This implies that $|\{\kappa-\Delta, \kappa-\Delta+1\} \cap \mathcal{U}(u_2)| = 1$, say $\kappa-\Delta \in \mathcal{U}(u_2)$. Hence, we have $\varepsilon_i = \kappa-\Delta$ and $\mathcal{A}(u_2) \subseteq \mathcal{U}(u_3)$.

Suppose that there exists no $(2, 1, u, w)$-critical path. Thus, there exists a $(\mu_i, \alpha_i, w, w_1)$-critical path with $\mu_i \in \mathcal{U}(w) \setminus \{1, 2, 3\}$; otherwise, reassigning α_i to ww_1 and 1 to wu will result in an acyclic edge coloring of G. Note that $\mathcal{U}(w_1) \supseteq \{1, 3, \kappa-\Delta, \kappa-\Delta+1, \mu_1\} \cup \mathcal{A}(u_2)$, it follows that $\mathcal{U}(u_2) \cap (\mathcal{U}(u) \cup \mathcal{U}(w)) = \{2, \kappa-\Delta\}$ and $|\mathcal{U}(u_2) \cap C(wu)| = \Delta-2$. Moreover, $\mathcal{U}(w_1) = \{1, 3, \kappa-\Delta, \kappa-\Delta+1, \mu_1\} \cup \mathcal{A}(u_2)$ and $|\mathcal{A}(u_2)| = 1$, say $\mu_1 = 4$. Thus, there exists a $(\kappa-\Delta, 1, u, u_2)$-critical path; otherwise, reassigning 1 to uu_2 will take us back to Subcase 2.1.1. So, we have $1 \in \mathcal{U}(u_3)$. Furthermore, there exists a $(\kappa-\Delta, 4, u, u_2)$-critical path; otherwise, reassigning 4 to uu_2 and α_1 to wu results in an acyclic edge coloring of G. Hence, $\{1, 4, \kappa-\Delta\} \subseteq \mathcal{U}(u_3)$. Recall that $|\mathcal{A}(u_3)| \geq 2$ and $|\mathcal{A}(u_2)| = 1$, it follows that $\mathcal{A}(w_1) \cap \mathcal{A}(u_3) \neq \emptyset$, say $\alpha_1 \notin \mathcal{U}(u_3)$. If $1 \notin \mathcal{U}(u_4)$, then reassigning 1 to uu_4 and α_1 to uw_1 will take us back to Subcase 2.1.2. Thus, we have $1 \in \mathcal{U}(u_4)$. If $2 \notin \mathbb{S}$, then reassigning $2, \beta_1$ and α_1 to uu_3, uu_2 and uw respectively results in an acyclic edge coloring of G. Thus $2 \in \mathbb{S}$. If $3 \notin \mathbb{S}$, then reassigning $3, \alpha_1$ and β_1 to uu_4, uw_1 and uw respectively results in an acyclic edge coloring of G. Thus $3 \in \mathbb{S}$. If $5 \notin \mathbb{S}$, then there exists a $(5, \alpha_1, w, u_2)$-alternating path, otherwise, reassigning 5 to

uu_2 and α_1 to uw results in an acyclic edge coloring of G; but reassigning $5, \beta_1$ and α_1 to uu_3, uu_2 and uw results in an acyclic edge coloring of G. Thus $5 \in \mathbb{S}$. Similarly, $\{5, 6, \ldots, \deg_G(w) - 1\} \subseteq \mathbb{S}$. Therefore, we have

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq 3|\{1\}| + 2|\{4, \kappa - \Delta\}| + |\{2, 3, \kappa - \Delta + 1, 5, 6, \ldots, \deg_G(w) - 1\}| = \deg_G(w) + 5.$$

Suppose that there exists a $(2, 1, w, u)$-critical path. It follows that $\{1, 2, \kappa - \Delta\} \subseteq \mathcal{U}(u_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u))$. It is obvious that $\mathcal{A}(u_2) \subseteq \mathcal{U}(w_1)$, thus $\mathcal{U}(w_1) = \{1, 3, \kappa - \Delta, \kappa - \Delta + 1\} \cup \mathcal{A}(u_2)$, $\mathcal{U}(u_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, 2, \kappa - \Delta\}$ and $|\mathcal{U}(u_2) \cap C(wu)| = \Delta - 3$. If $\mathcal{A}(w_1) \subseteq \mathcal{U}(u_3)$, then $\mathcal{U}(u_3) = \mathcal{A}(w_1) \cup \mathcal{A}(u_2) \cup \{\kappa - \Delta\} = C(wu) \cup \{\kappa - \Delta\}$; but reassigning $2, \beta_1$ and α_1 to uu_3, uu_2 and uw respectively results in an acyclic edge coloring of G. So we may assume that $\mathcal{A}(w_1) \nsubseteq \mathcal{U}(u_3)$ and $\alpha_1 \notin \mathcal{U}(u_3)$. If $3 \notin \mathbb{S}$, then reassigning $3, \alpha_1$ and β_1 to uu_4, uw_1 and uw respectively results in an acyclic edge coloring of G. Thus, we have $3 \in \mathbb{S}$. For every color θ in $\mathcal{U}(w) \setminus \{3\}$, we have that $\theta \in \mathbb{S}$; otherwise, reassigning θ, β_1 and α_1 to uu_3, uu_2 and uw respectively results in an acyclic edge coloring of G. If $1 \notin \mathcal{U}(u_3) \cup \mathcal{U}(u_4)$, then reassigning 1 to uu_4 and α_1 to uw_1 will take us back to Subcase 2.1.1. Hence, the color 1 appears exactly three times in $\mathbb{S}$. If $\kappa - \Delta + 1$ appears at least twice in $\mathbb{S}$ or $|X| \geq 1$, then

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq \deg_G(w) + 5.$$

So we may assume that $\kappa - \Delta + 1$ appears precisely once (at w_1) and $X = \emptyset$. Note that $\beta_1 \notin \mathcal{U}(u_4)$. But reassigning β_1 to uu_4 and $\kappa - \Delta + 1$ to uu_2 will take us back to Case 1.

Subcase 2.1.3.2. Now, we may assume that $\{\kappa - \Delta, \kappa - \Delta + 1\} \nsubseteq \mathcal{U}(w_1)$ and $\kappa - \Delta + 1 \notin \mathcal{U}(w_1)$.

Thus, there exists a $(\kappa - \Delta, \alpha_i, u, w_1)$-critical path for every α_i; otherwise, reassigning α_i to uw_1 will take us back to Case 1. It follows that $\kappa - \Delta \in \mathcal{U}(w_1)$ and $\mathcal{A}(w_1) \subseteq \mathcal{U}(u_3) \cap \mathcal{U}(u_2)$. If $\Upsilon(uu_3) \subseteq C(uw)$, then reassigning α_1 to uw_1 and 1 to uu_3 will take us back to Subcase 2.1.2. So we may assume that $\Upsilon(uu_3) \nsubseteq C(wu)$ and $C(wu) \nsubseteq \Upsilon(uu_3)$.

(1) Suppose that $\mathcal{A}(u_2) \cap \mathcal{A}(u_3) = \emptyset$. It follows that $\mathcal{A}(w_1), \mathcal{A}(u_2)$ and $\mathcal{A}(u_3)$ are pairwise disjoint. Suppose that there exists no $(2, 1, u, w)$-critical path. Thus, there exists a (τ, α_1, w, w_1)-critical path, where $\tau \in \mathcal{U}(w) \setminus \{1, 2, 3\}$; otherwise, reassigning 1 to uw and α_1 to ww_1 results in an acyclic edge coloring of G. Since $\mathcal{U}(u_3) \supseteq \mathcal{A}(w_1) \cup \mathcal{A}(u_2)$ and $C(wu) \nsubseteq \mathcal{U}(u_3)$, it follows that $|\mathcal{A}(w_1)| = \Delta - 3$, $|\mathcal{A}(u_2)| = 1$ and $|\mathcal{U}(u_3) \cap (\mathcal{U}(w) \cup \mathcal{U}(u))| = 2$. If $1 \notin \mathcal{U}(u_3)$, then there exists a $(\kappa - \Delta + 1, 1, u, u_3)$-critical path; otherwise, reassigning 1 to uu_3 and α_1 to uw_1 will take us back to Subcase 2.1.2. Thus, $\Upsilon(uu_3) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1\}$ or $\{\kappa - \Delta + 1\}$. If there exists no $(\kappa - \Delta + 1, 3, u, u_3)$-critical path, then reassigning $3, \alpha_1$ and β_1 to uu_3, uw_1 and uw results in an acyclic edge coloring of G. Hence, there exists a $(\kappa - \Delta + 1, 3, u, u_3)$-critical path and $\Upsilon(uu_3) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{\kappa - \Delta + 1\}$. But reassigning 1 to uu_2 will take us back to Subcase 2.1.1.

Now, we consider the other subcase: suppose that there exists a $(2, 1, u, w)$-critical path and $1 \in \mathcal{U}(u_2)$. Since $\mathcal{U}(u_3) \supseteq \mathcal{A}(w_1) \cup \mathcal{A}(u_2)$ and $C(wu) \nsubseteq \mathcal{U}(u_3)$, so we have that $|\mathcal{A}(w_1)| = \Delta - 4$, $|\mathcal{A}(u_2)| = 2$ and $\mathcal{U}(w_1) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, 3, \kappa - \Delta\}$, $\mathcal{U}(u_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, 2, \varepsilon_1\}$ and $|\mathcal{U}(u_3) \cap (\mathcal{U}(w) \cup \mathcal{U}(u))| = 2$. If $1 \in \mathcal{U}(u_3)$, then $\mathcal{U}(u_3) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, \kappa - \Delta\}$, and then reassigning β_1, α_1 and 3 to uw, uw_1 and uu_3 results in an cyclic edge coloring. Thus, $1 \notin \mathcal{U}(u_3)$. There exists a $(\kappa - \Delta + 1, 1, u, u_3)$-critical path; otherwise, reassigning 1 to uu_3 and α_1 to uw_1 will take us back to Subcase 2.1.2. This implies that $\mathcal{U}(u_3) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{\kappa - \Delta, \kappa - \Delta + 1\}$ and 1 appears three times in $\mathbb{S}$. There exists a $(\kappa - \Delta + 1, 3, u, u_3)$-critical path, otherwise, reassigning β_1, α_1 and 3 to uw, uw_1 and uu_3 results in an acyclic edge coloring of G. Now, we have $\{1, 3\} \subseteq \Upsilon(uu_4)$. If $\beta \in \mathcal{A}(u_2) \cap \mathcal{A}(u_4)$, then $\varepsilon_1 = \kappa - \Delta$ and there exists a $(\kappa - \Delta, \beta, u, u_2)$-critical path; but reassigning β to uu_4 and $\kappa - \Delta + 1$ to uw results in an acyclic edge coloring of G. This implies that $\mathcal{A}(u_2) \subseteq \mathcal{U}(u_4)$ and $\mathcal{A}(u_2) \subseteq X$. Suppose that $\{4, 5, \ldots, \deg_G(w) - 1\} \nsubseteq \mathbb{S}$. So, by symmetry, we may assume that $4 \notin \mathbb{S}$. There exists a $(4, \beta_1, w, w_1)$-alternating path; otherwise, reassigning 4 to uw_1 and β_1 to uw results in an acyclic edge coloring of G. But reassigning $4, \alpha_1$ and β_1 to uu_3, uw_1 and uw results in an acyclic edge coloring of G. Hence, $\{3, 4, \ldots, \deg_G(w) - 1\} \subseteq \mathbb{S}$.

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq |\{3, 4, \ldots, \deg_G(w) - 1\}| + 3|\{1\}| + |\{\varepsilon_1\}| + |\{\kappa - \Delta\}| + |\{\kappa - \Delta + 1\}| + |\mathcal{A}(u_2)| \geq \deg_G(w) + 5.$$

(2) So we may assume that $\mathcal{A}(u_2) \cap \mathcal{A}(u_3) \neq \emptyset$, say $\beta_1 \in \mathcal{A}(u_2) \cap \mathcal{A}(u_3)$. Thus, there exists a $(\kappa - \Delta + 1, \beta_1, u, u_2)$-critical path; otherwise, reassigning β_1 to uu_2 will take us back to Case 1. So, we have $\kappa - \Delta + 1 \in \mathcal{U}(u_2)$.

Suppose that the color 1 only appears once (at w_1) in $\mathbb{S}$. If there exists no $(3,1,u,u_2)$-critical path, then reassigning 1 to uu_2 will take us back to Subcase 2.1.1. But if there exists a $(3,1,u,u_2)$-critical path, then reassigning 1 to uu_4 and β_1 to uu_2 will take us back to Subcase 2.1.1 again. Hence, the color 1 appears at least twice in $\mathbb{S}$.

If $2 \notin \mathbb{S}$, then reassigning $2, \beta_1$ and α_1 to uu_4, uu_2 and uw respectively results in an acyclic edge coloring of G. If $3 \notin \mathbb{S}$, then reassigning $3, \alpha_1$ and β_1 to uu_3, uw_1 and uw respectively, results in an acyclic edge coloring of G. Suppose that $4 \notin \mathbb{S}$. There exists a $(4, \beta_1, w, w_1)$-alternating path; otherwise, reassigning 4 to uw_1 and β_1 to uw results in an acyclic edge coloring of G. Now, reassigning $4, \alpha_1$ and β_1 to uu_3, uw_1 and uw respectively results in an acyclic edge coloring of G. Thus, $\{2,3,4\} \subseteq \mathbb{S}$. By symmetry, we have that $\mathcal{U}(w) \setminus \{1,2,3\} \subseteq \mathbb{S}$.

Suppose that $\kappa - \Delta$ appears only once (at w_1) in $\mathbb{S}$. Thus, there exists a $(3, \kappa-\Delta, u, w)$-critical path; otherwise, reassigning $\kappa - \Delta$ to uw and β_1 to uu_3 results in an acyclic edge coloring of G. But reassigning β_1 to uu_3 and $\kappa - \Delta$ to uu_2 will take us back to Case 1. Hence, the color $\kappa - \Delta$ appears at least twice in $\mathbb{S}$.

Note that $|\mathcal{A}(w_1)| \geq 2$. If $\mathcal{A}(w_1) \subseteq \mathcal{U}(u_4)$, then $\mathcal{A}(w_1) \subseteq X$, and then

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq |\{\kappa - \Delta + 1, 2, 3, \ldots, \deg_G(w) - 1\}| + 2|\{1, \kappa - \Delta\}| + |\mathcal{A}(w_1)| \geq \deg_G(w) + 5.$$

So we may assume that $\mathcal{A}(w_1) \nsubseteq \mathcal{U}(u_4)$, say $\alpha_1 \notin \mathcal{U}(u_4)$. There exists a $(2, \kappa - \Delta + 1, w, u)$-critical path; otherwise, reassigning α_1 to uu_4 and $\kappa - \Delta + 1$ to uw results in an acyclic edge coloring of G. Consequently, there exists a $(\kappa - \Delta, \kappa - \Delta + 1, u, w_1)$-critical path and $\kappa - \Delta + 1 \in \mathcal{U}(u_3)$; otherwise, reassigning α_1 to uu_4 and $\kappa - \Delta + 1$ to uw_1 will take us back to Case 1. Hence, the color $\kappa - \Delta + 1$ appears exactly twice in $\mathbb{S}$.

Suppose that there exists no $(2,1,u,w)$-critical path. Thus, there exists a (τ, α_1, w, w_1)-critical path with $\tau \in \mathcal{U}(w) \setminus \{1,2,3\}$; otherwise, reassigning 1 to uw and α_1 to ww_1 results in an acyclic edge coloring of G. If τ only appears once (at w_1) in $\mathbb{S}$, then reassigning τ, α_1 and β_1 to uu_3, uw_1 and uw respectively results in an acyclic edge coloring of G. Hence, the color τ appears at least twice in $\mathbb{S}$. Hence,

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq |\{2, 3, \ldots, \deg_G(w) - 1\}| + 2|\{1, \kappa - \Delta, \kappa - \Delta + 1\}| + |\{\tau\}| = \deg_G(w) + 5.$$

Suppose there exists a $(2,1,u,w)$-critical path and $1 \in \mathcal{U}(u_2)$. If $1 \notin \mathcal{U}(u_3) \cup \mathcal{U}(u_4)$, then reassigning 1 to uu_3 and α_1 to uw_1 will take us back to Subcase 2.1.1. Hence, the color 1 appears at least three times in $\mathbb{S}$,

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq |\{2, 3, \ldots, \deg_G(w) - 1\}| + 3|\{1\}| + 2|\{\kappa - \Delta, \kappa - \Delta + 1\}| = \deg_G(w) + 5.$$

Subcase 2.1.4. Neither the color on w_1w nor the color on w_1u is a common color.

By symmetry, assume that $\phi(uw_1) = \kappa - \Delta$, $\phi(uu_2) = 2$, $\phi(uu_3) = 3$ and $\phi(uu_4) = \kappa - \Delta + 1$.

If $\Upsilon(uu_2) \subseteq C(wu)$, then reassigning 1 to uu_2 will take us back to Subcase 2.1.2. This implies that $\Upsilon(uu_2) \nsubseteq C(wu)$ and $\mathcal{A}(u_2) \neq \emptyset$. Thus, there exists a $(\varepsilon_i, \beta_i, u, u_2)$-critical path with $\varepsilon_i \in \{\kappa - \Delta, \kappa - \Delta + 1\}$; otherwise, reassigning β_i to uu_2 will take us back to Case 1. Similarly, we have that $\Upsilon(uu_3) \nsubseteq C(wu)$ and $\mathcal{A}(u_3) \neq \emptyset$, and thus there exists a (m_i, ξ_i, u, u_3)-critical path with $m_i \in \{\kappa - \Delta, \kappa - \Delta + 1\}$. If $1 \notin \mathcal{U}(u_2) \cup \mathcal{U}(u_3)$, then reassigning 1 to uu_2 will create a $(1, \kappa - \Delta + 1)$-dichromatic cycle containing uu_2, otherwise, it will take us back to Subcase 2.1.2; but reassigning 1 to uu_3 will take us back to Subcase 2.1.2 again. It follows that $1 \in \mathcal{U}(u_2) \cup \mathcal{U}(u_3)$ and 1 appears at least twice in $\mathbb{S}$.

Subcase 2.1.4.1. Suppose that $\mathcal{A}(u_2) \cup \mathcal{A}(u_3) \nsubseteq \mathcal{U}(w_1)$ and $\beta_1 = \alpha_1 \notin \mathcal{U}(w_1)$.

Hence, there exists a $(3, \beta_1, u, w)$-critical path and $(\kappa - \Delta + 1, \beta_1, u, u_2)$-critical path, thus $\kappa - \Delta + 1 \in \mathcal{U}(u_2)$.

There exists a $(2, \kappa - \Delta, u, w)$- or $(3, \kappa - \Delta, u, w)$-critical path; otherwise, reassigning β_1 to uw_1 and $\kappa - \Delta$ to uw results in an acyclic edge coloring of G. It follows that $\kappa - \Delta \in \mathcal{U}(u_2) \cup \mathcal{U}(u_3)$. Moreover, $\kappa - \Delta$ appears at least twice in $\mathbb{S}$; otherwise, assume that $\kappa - \Delta$ only appears at u_2, thus reassigning $\kappa - \Delta$ to uu_3 and β_1 to uw_1 will take us back to Case 1.

If $2 \notin \mathbb{S}$, then reassigning $\beta_1, 2$ and ξ_1 to uu_2, uu_4 and uw respectively results in an acyclic edge coloring of G. Thus $2 \in \mathbb{S}$.

Suppose that $4 \notin \mathbb{S}$. There exists a $(4, \xi_1, w, u_2)$-alternating path for every $\xi_i \in \mathcal{A}(u_3)$; otherwise, reassigning 4 to uu_2 and ξ_1 to uw results in an acyclic edge coloring of G. Now, reassigning $\beta_1, 4$ and ξ_1 to uu_2, uu_4 and uw respectively results in an acyclic edge coloring of G again. Hence, the color 4 appears in $\mathbb{S}$. Similarly, we can prove that $\mathcal{U}(w) \setminus \{1,2,3\} \subseteq \mathbb{S}$.

Suppose that $3 \notin \mathbb{S}$. If there exists no $(\kappa - \Delta + 1, \xi_i, u, u_3)$-critical path, then reassigning 3 to uw_1 and ξ_i to uu_3 will take us back to Subcase 2.1.3. Hence, there exists a $(\kappa - \Delta + 1, \xi_i, u, u_3)$-critical path for every $\xi_i \in \mathcal{A}(u_3)$,

and then $\kappa-\Delta+1 \in \mathcal{U}(u_3)$ and $\mathcal{A}(u_3) \subseteq \mathcal{U}(u_4)$. If there exists no $(\kappa-\Delta, \xi_i, u, u_3)$-critical path, then reassigning $3, \xi_i$ and β_1 to uu_4, uu_3 and uw respectively, results in an acyclic edge coloring of G. Hence, both $(\kappa-\Delta, \xi_i, u, u_3)$- and $(\kappa-\Delta+1, \xi_i, u, u_3)$-critical path exist for every $\xi_i \in \mathcal{A}(u_3)$, and then $\{\kappa-\Delta, \kappa-\Delta+1\} \subseteq \mathcal{U}(u_3)$ and $\mathcal{A}(u_3) \subseteq \mathcal{U}(w_1) \cap \mathcal{U}(u_4)$. Clearly, every color in $\mathcal{A}(u_3)$ appears precisely three times in $\mathbb{S}$. Therefore,

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq |\{2\} \cup \{4, \ldots, \deg_G(w)-1\}| + 2|\{1, \kappa-\Delta, \kappa-\Delta+1\}| + |\mathcal{A}(u_3)| \geq \deg_G(w)+5.$$

So, in the following, we may assume that $3 \in \mathbb{S}$.

If there exists a $(2, 1, u, w)$-critical path (or $(3, 1, u, w)$-critical path) and 1 appears only twice in $\mathbb{S}$, then reassigning 1 to uu_3 (to uu_2) will take us back to Subcase 2.1.2. In other words, if there exists a $(2, 1, u, w)$-critical path or $(3, 1, u, w)$-critical path, then the color 1 appears at least three times in $\mathbb{S}$.

Suppose that neither $(2, 1, u, w)$-critical path nor $(3, 1, u, w)$-critical path exists. If there exists no (τ, β_1, w, w_1)-critical path with some $\tau \in \mathcal{U}(w) \setminus \{1, 3\}$, then reassigning 1 to uw and β_1 to ww_1 results in an acyclic edge coloring of G. Hence, there exists a (τ, β_1, w, w_1)-critical path with some $\tau \in \mathcal{U}(w) \setminus \{1, 3\}$. Suppose that there exists a $(2, \beta_1, w, w_1)$-critical path and 2 appears only once in $\mathbb{S}$. This implies that there exists a $(\kappa-\Delta, 2, u, u_4)$-critical path; otherwise reassigning $2, \beta_1$ and ξ_1 to uu_4, uu_2 and uw respectively results in an acyclic edge coloring of G. But reassigning $2, \beta_1, \xi_1$ and ζ^* ($\zeta^* = \beta_2$ if $|\mathcal{A}(u_2)| \geq 2$, otherwise, $\zeta^* = \kappa-\Delta$) to uu_4, uw_1, uw and uu_2 respectively, and we obtain an acyclic edge coloring of G. Thus, if there exists a $(2, \beta_1, w, w_1)$-critical path, then the color 2 appears at least twice in $\mathbb{S}$. Suppose that there exists a $(4, \beta_1, w, w_1)$-critical path and 4 only appears once in $\mathbb{S}$. Hence, there is a $(\kappa-\Delta, 4, u, u_3)$-critical path, otherwise, reassigning 4 to uu_3 and β_1 to uw results in an acyclic edge coloring of G. Now, reassigning $4, \beta_1$ and ξ_1 to uu_4, uu_2 and uw will create a $(4, \xi_1)$-dichromatic cycle containing uw; otherwise, the resulting coloring is an acyclic edge coloring of G. But reassigning 4 to uu_2 and ξ_1 to uw results in an acyclic edge coloring of G. Thus, if there exists a $(4, \beta_1, w, w_1)$-critical path, then the color 4 appears at least twice in $\mathbb{S}$. Similarly, if there exists a (τ, β_1, w, w_1)-critical path with $\tau \geq 4$, then the color τ appears at least twice in $\mathbb{S}$. Therefore, the color τ appears at least twice in $\mathbb{S}$.

By the above arguments, regardless of the existence of $(2, 1, u, w)$-critical path or $(3, 1, u, w)$-critical path, if $\kappa-\Delta+1$ appears at least twice or $|X| \geq 1$, then

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq \deg_G(w)+5.$$

So we may assume that the color $\kappa-\Delta+1$ appears only once (at u_2) in $\mathbb{S}$ and $X = \emptyset$. If $\mathcal{A}(u_3) \nsubseteq \mathcal{U}(w_1)$, say $\xi_1 \notin \mathcal{U}(w_1)$, then there exists a $(2, \xi_1, u, w)$-critical path and $(\kappa-\Delta+1, \xi_1, u, u_3)$-critical path, and then $\kappa-\Delta+1 \in \mathcal{U}(u_3)$, a contradiction. So we may assume that $\mathcal{A}(u_3) \subseteq \mathcal{U}(w_1)$ and $\mathcal{A}(u_3) \cap \mathcal{U}(u_4) = \emptyset$.

Clearly, there exists a $(2, \xi_1, u, w)$-critical path. Thus, there exists a $(\kappa-\Delta, \xi_1, u, u_3)$-critical path; otherwise, reassigning ξ_1 to uu_3 will take us back to Case 1. Hence, there exists a $(2, \kappa-\Delta+1, u, w)$-critical path; otherwise, reassigning ξ_1 to uu_4 and $\kappa-\Delta+1$ to uw will result in an acyclic edge coloring of G. Now, reassigning ξ_1 to uu_4 and $\kappa-\Delta+1$ to uu_3 will take us back to Case 1.

Subcase 2.1.4.2. $\mathcal{A}(u_2) \cup \mathcal{A}(u_3) \subseteq \mathcal{U}(w_1)$.

Firstly, suppose that $\mathcal{A}(u_2) \cup \mathcal{A}(u_3) \nsubseteq \mathcal{U}(u_4)$ and $\beta_1 = \zeta_1 \notin \mathcal{U}(u_4)$. Hence, there exists a $(3, \beta_1, u, w)$-critical path and a $(\kappa-\Delta, \beta_1, u, u_2)$-critical path, and then $\kappa-\Delta \in \mathcal{U}(u_2)$.

If $\{2, 3, \kappa-\Delta+1\} \cap \mathcal{U}(w_1) = \emptyset$, then reassigning β_1 to uu_2 and 2 to uw_1 will take us back to Subcase 2.1.3. Hence, $\{2, 3, \kappa-\Delta+1\} \cap \mathcal{U}(w_1) \neq \emptyset$. Recall that $1 \in \mathcal{U}(u_2) \cup \mathcal{U}(u_3)$. Since $\mathcal{A}(u_2) \cup \mathcal{A}(u_3) \subseteq \mathcal{U}(w_1)$, it follows that $\deg_G(w_1) = 6$, $\deg_G(w) = \kappa-\Delta$, and $|\mathcal{A}(u_2)| + |\mathcal{A}(u_3)| = 3$. Furthermore, we have that $\{\kappa-\Delta, \kappa-\Delta+1\} \cap \mathcal{U}(u_2) = \{\kappa-\Delta\}$ and $|\{\kappa-\Delta, \kappa-\Delta+1\} \cap \mathcal{U}(u_3)| = 1$. Thus, there exists a $(3, \kappa-\Delta+1, w, u)$-critical path; otherwise reassigning β_1 to uu_4 and $\kappa-\Delta+1$ to uw will result in an acyclic edge coloring of G. Hence, $\{\kappa-\Delta, \kappa-\Delta+1\} \cap \mathcal{U}(u_3) = \{\kappa-\Delta+1\}$.

If $1 \notin \mathcal{U}(u_2)$, then $\mathcal{U}(u_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{2, \kappa-\Delta\}$, but reassigning 1 to uu_2 will take us back to Subcase 2.1.2. This implies that $\mathcal{U}(u_2) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, 2, \kappa-\Delta\}$ and $\mathcal{U}(u_3) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{3, \kappa-\Delta+1\}$. Now, there is a $(\kappa-\Delta+1, 1, u, u_3)$-critical path; otherwise, reassigning 1 to uu_3 will take us back to Subcase 2.1.2. Thus, there exists a $(\kappa-\Delta, \kappa-\Delta+1, u, u_2)$-critical path; otherwise, reassigning β_1 to uu_4 and $\kappa-\Delta+1$ to uu_2 will take us back to Case 1. Hence, $\mathcal{U}(w_1) \cap (\mathcal{U}(w) \cup \mathcal{U}(u)) = \{1, \kappa-\Delta, \kappa-\Delta+1\}$. Moreover, there exists a $(\kappa-\Delta+1, 2, u, w_1)$-critical path; otherwise, reassigning $\beta_1, 2$ and $\kappa-\Delta$ to uu_2, uw_1 and uw respectively, results in an acyclic edge coloring of G. It is obvious that $2 \in \mathcal{U}(u_4)$. If $\kappa-\Delta \notin \mathcal{U}(u_4)$, then reassigning $\kappa-\Delta, \beta_1, 2$ and ξ_1 to uu_4, uu_2, uw_1 and uw respectively,

results in an acyclic edge coloring of G. Thus, $\kappa-\Delta\in\mathcal{U}(u_4)$. Recall that $\{1,2\}\subseteq\mathcal{U}(u_4)$. If there exists a color τ in $\mathcal{U}(w)\setminus\mathcal{U}(u_4)$, then reassigning τ, ξ_1 and β_1 to uu_4, uu_3 and uw respectively, will result in an acyclic edge coloring of G. Hence, $\mathcal{U}(w)\subseteq\mathcal{U}(u_4)$. Then

$$\sum_{\theta\in\mathcal{U}(w)\cup\mathcal{U}(u)}\mathrm{mul}_{\mathbb{S}}(\theta)+|X|\geq|\mathcal{U}(w)|+2|\{1,\kappa-\Delta,\kappa-\Delta+1\}|=\deg_G(w)+5.$$

Secondly, suppose that $\mathcal{A}(u_2)\cup\mathcal{A}(u_3)\subseteq\mathcal{U}(u_4)$. Thus, every color in $\mathcal{A}(u_2)\cup\mathcal{A}(u_3)$ appears three times in $\mathbb{S}$, and then $\mathcal{A}(u_2)\cup\mathcal{A}(u_3)\subseteq X$. Recall that $1\in\mathcal{U}(u_2)\cup\mathcal{U}(u_3)$. Since $\mathcal{A}(u_2)\cup\mathcal{A}(u_3)\subseteq\mathcal{U}(w_1)$, it follows that $|\{\kappa-\Delta,\kappa-\Delta+1\}\cap\mathcal{U}(u_2)|=1$ or $|\{\kappa-\Delta,\kappa-\Delta+1\}\cap\mathcal{U}(u_3)|=1$.

Suppose that $1\in\Upsilon(uu_2)\cap\Upsilon(uu_3)$. It follows that $\deg_G(w)=6$ and $\mathcal{U}(w_1)=\{1,\kappa-\Delta\}\cup\mathcal{A}(u_2)\cup\mathcal{A}(u_3)$. Moreover, we have that $|\{\kappa-\Delta,\kappa-\Delta+1\}\cap\Upsilon(uu_2)|=1$ and $|\{\kappa-\Delta,\kappa-\Delta+1\}\cap\Upsilon(uu_3)|=1$. If $\kappa-\Delta+1\notin\Upsilon(uu_2)$, then reassigning $\beta_1,2$ and ξ_1 to uu_2, uw_1 and wu respectively results in an acyclic edge coloring of G. So we have that $\kappa-\Delta+1\in\Upsilon(uu_2)$. Similarly, we have that $\kappa-\Delta+1\in\Upsilon(uu_3)$. But reassigning α_1 to uw_1 and $\kappa-\Delta$ to wu results in an acyclic edge coloring of G.

So we may assume that $1\notin\Upsilon(uu_2)\cap\Upsilon(uu_3)$ and $1\in\Upsilon(uu_2)$. If there is a $(2,1,u,u_3)$-critical path, then $\mathcal{U}(w_1)=\{1,\kappa-\Delta\}\cup\mathcal{A}(u_2)\cup\mathcal{A}(u_3)$ and $2\in\mathcal{U}(u_3)$, but reassigning α_1 to ww_1 and 1 to wu results in an acyclic edge coloring of G. Hence, there exists no $(2,1,u,u_3)$-critical path. Thus, there exists a $(\kappa-\Delta+1,1,u,u_3)$-critical path, otherwise, reassigning 1 to uu_3 will take us back to Subcase 2.1.2. This implies that the color 1 appears at least three times in $\mathbb{S}$.

Suppose that $3\notin\mathbb{S}$. Thus, there exists a $(2,\kappa-\Delta+1,w,u)$-critical path; otherwise, reassigning $3,1$ and $\kappa-\Delta+1$ to uu_4, uu_3 and uw respectively, results in an acyclic edge coloring of G. Hence, $\kappa-\Delta+1\in\mathcal{U}(u_2)\cap\mathcal{U}(u_3)$. If $\kappa-\Delta\notin\mathcal{U}(u_3)$, then reassigning $3,\xi_1$ and β_1 to uu_4, uu_3 and uw respectively, results in an acyclic edge coloring of G. So we may assume that $\kappa-\Delta\in\mathcal{U}(u_3)$. Hence, $|\mathcal{A}(u_2)|=|\mathcal{A}(u_3)|=2$ and $\mathcal{U}(w_1)=\{1,\kappa-\Delta\}\cup\mathcal{A}(u_2)\cup\mathcal{U}(u_3)$. Now, reassigning $3,1,\beta_1$ and α_1 to uu_4, uu_3, uw and ww_1, results in an acyclic edge coloring of G. Therefore, we can conclude that $3\in\mathbb{S}$.

Suppose that $4\notin\mathbb{S}$. Thus, there is a $(4,\beta_1,w,u_3)$-alternating path; otherwise, reassigning 4 to uu_3 and β_1 to uw will result in an acyclic edge coloring of G. Similarly, there exists a $(4,\xi_1,w,u_2)$-alternating path. Moreover, there exists a $(\kappa-\Delta,\xi_1,u,u_3)$-critical path; otherwise, reassigning 4, ξ_1 and β_1 to uu_4, uu_3 and uw respectively, results in an acyclic edge coloring of G. Thus, $\mathcal{U}(u_3)\cap(\mathcal{U}(w)\cup\mathcal{U}(u))=\{3,\kappa-\Delta,\kappa-\Delta+1\}$, $|\mathcal{A}(u_3)|=2$ and $|\mathcal{A}(u_2)|=2$. Hence, $|\mathcal{U}(u_2)\cap\{\kappa-\Delta,\kappa-\Delta+1\}|=1$. If $\kappa-\Delta\notin\mathcal{U}(u_2)$, then reassigning $4,\beta_1$ and ξ_1 to uu_4, uu_2 and uw respectively, results in an acyclic edge coloring of G. Hence, we have that $\mathcal{U}(u_2)\cap(\mathcal{U}(w)\cup(u))=\{1,2,\kappa-\Delta\}$. But reassigning $\xi_1,4$ and β_1 to uw, uw_1 and uu_2 respectively, results in an acyclic edge coloring of G. So, $4\in\mathbb{S}$. Similarly, we have that $\mathcal{U}(w)\setminus\{1,2,3\}\subseteq\mathbb{S}$.

Recall that $|\mathcal{A}(u_2)\cup\mathcal{A}(u_3)|\geq3$, $|\{\kappa-\Delta,\kappa-\Delta+1\}\cap\mathcal{U}(u_2)|\geq1$ and $|\{\kappa-\Delta,\kappa-\Delta+1\}\cap\mathcal{U}(u_3)|\geq1$. Hence,

$$\sum_{\theta\in\mathcal{U}(w)\cup\mathcal{U}(u)}\mathrm{mul}_{\mathbb{S}}(\theta)+|X|\geq|\{3,4,\ldots,\deg_G(w)-1\}|+3|\{1\}|+1+1+|\mathcal{A}(u_2)\cup\mathcal{A}(u_3)|\geq\deg_G(w)+5.$$

Subcase 2.2. $\mathcal{U}(w)\cap\mathcal{U}(u)=\{\lambda_1,\lambda_2,\lambda_3\}$ and $w_1=u_1$. Note that $|C(wu)|\geq\Delta$.

Subcase 2.2.1. The color on uw_1 is a common color.

By symmetry, assume that $\phi(uw_1)=\lambda_1$, $\phi(uu_2)=\lambda_2$, $\phi(uu_3)=\lambda_3$ and $\phi(uu_4)=\kappa-\Delta$.

If $\Upsilon(uu_2)\subseteq C(wu)$, then reassigning β_1 to uu_2 will take us back to Subcase 2.1. So we have that $\Upsilon(uu_2)\nsubseteq C(wu)$ and $|\mathcal{U}(u_2)\cap C(wu)|\leq\Delta-2$; similarly, we also have that $|\mathcal{U}(u_3)\cap C(wu)|\leq\Delta-2$. If $\mathcal{U}(w_1)\cap(\mathcal{U}(w)\cup\mathcal{U}(u))=\{1,\lambda_1\}$, then reassigning α_1 to uw_1 will take us back to Subcase 2.1 again. Hence, $|\mathcal{U}(w_1)\cap(\mathcal{U}(w)\cup\mathcal{U}(u))|\geq3$. By Claim 2, we have $\mathcal{A}(u_2)\cup\mathcal{A}(u_3)\subseteq\mathcal{U}(w_1)$ and $\mathcal{A}(u_2)\cap\mathcal{A}(u_3)=\emptyset$. Further, we have that $|\mathcal{U}(w_1)|\geq3+|\mathcal{A}(u_2)|+|\mathcal{A}(u_3)|>\deg_G(w_1)$, which is a contradiction.

Subcase 2.2.2. The color on uw_1 is not a common color, but the color on ww_1 is a common color.

By symmetry, assume that $\phi(uw_1)=\kappa-\Delta$, $\phi(uu_2)=2$, $\phi(uu_3)=3$ and $\phi(uu_4)=1$.

If $\Upsilon(uu_2)\subseteq C(wu)$, then reassigning β_1 to uu_2 will take us back to Subcase 2.1. So we have that $\Upsilon(uu_2)\nsubseteq C(wu)$ and $|\mathcal{U}(u_2)\cap C(wu)|\leq\Delta-2$; similarly, we also have that $|\mathcal{U}(u_3)\cap C(wu)|\leq\Delta-2$ and $|\mathcal{U}(u_4)\cap C(wu)|\leq\Delta-2$.

If $\mathcal{U}(w_1)\cap(\mathcal{U}(w)\cup\mathcal{U}(u))=\{1,\kappa-\Delta\}$, then reassigning α_1 to ww_1 will take us back to Subcase 2.1 again. Hence, $|\mathcal{U}(w_1)\cap(\mathcal{U}(w)\cup\mathcal{U}(u))|\geq3$.

Furthermore, we have that $\mathcal{A}(u_2) \cup \mathcal{A}(u_3) \nsubseteq \mathcal{U}(w_1)$. Otherwise, if $\deg_G(w) \leq \kappa - \Delta$, then $|\mathcal{U}(w_1)| \geq 3 + |\mathcal{A}(u_2)| + |\mathcal{A}(u_3)| \geq 3 + 2 + 2 > 6$; and if $\deg_G(w) \leq \kappa - \Delta - 1$, then $|\mathcal{U}(w_1)| \geq 3 + |\mathcal{A}(u_2)| + |\mathcal{A}(u_3)| \geq 3 + 3 + 3 > 7$. Without loss of generality, assume that $\beta_1 \notin \mathcal{U}(w_1)$. Since $\beta_1 \notin \mathcal{U}(w_1) \cup \mathcal{U}(u_2)$, it follows that there exists a $(3, \beta_1, u, w)$-critical path. There exists a $(1, \beta_1, u, u_2)$-critical path; otherwise, reassigning β_1 to uu_2 will take us back to Subcase 2.1. Hence, $1 \in \mathcal{U}(u_2)$ and 1 appears at least twice in $\mathbb{S}$.

There exists a $(2, \kappa - \Delta, u, w)$- or $(3, \kappa - \Delta, u, w)$-critical path; otherwise, reassigning $\kappa - \Delta$ to uw and β_1 to uw_1 will result in an acyclic edge coloring of G. If $\kappa - \Delta$ appears only once in $\mathbb{S}$, then reassigning β_1 to uw_1 and $\kappa - \Delta$ to uu_4 will take us back to Subcase 2.1. Hence, the color $\kappa - \Delta$ appears at least twice in $\mathbb{S}$.

Let $t \in \mathcal{U}(w) \setminus \{1, 3\}$. If $t \notin \mathbb{S}$, then reassigning β_1 to uu_2 and t to uu_4 will take us back to Subcase 2.1. Hence, we have that $\mathcal{U}(w) \setminus \{1, 3\} \subseteq \mathbb{S}$.

If $\mathcal{A}(u_3) \subseteq \mathcal{U}(w_1)$, then every color in $\mathcal{A}(u_3)$ appears precisely three times in $\mathbb{S}$, and then

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq |\{2, 4, 5, \ldots, \deg_G(w) - 1\}| + 2|\{1, \kappa - \Delta\}| + |\mathcal{A}(u_3)| \geq \deg_G(w) + 3.$$

So we may assume that $\mathcal{A}(u_3) \nsubseteq \mathcal{U}(w_1)$ and $\xi_1 \notin \mathcal{U}(w_1) \cup \mathcal{U}(u_3)$. Similar to above, we can prove that there exists a $(2, \xi_1, w, u)$- and $(1, \xi_1, u, u_3)$-critical path, and then 1 appears precisely three times in $\mathbb{S}$. If $3 \notin \mathbb{S}$, then reassigning 3 to uu_4 and ξ_1 to uu_3 will take us back to Subcase 2.1. Thus, the color 3 appears at least once in $\mathbb{S}$. Therefore, we have

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq |\{2, 3, \ldots, \deg_G(w) - 1\}| + 2|\{\kappa - \Delta\}| + 3|\{1\}| = \deg_G(w) + 3.$$

Subcase 2.2.3. Neither the color on w_1w nor the color on w_1u is a common color.

By symmetry, assume that $\phi(uw_1) = \kappa - \Delta$, $\phi(uu_2) = 2$, $\phi(uu_3) = 3$ and $\phi(uu_4) = 4$.

If $\Upsilon(uu_2) \subseteq C(wu)$, then reassigning β_1 to uu_2 will take us back to Subcase 2.1. So we have that $\Upsilon(uu_2) \nsubseteq C(wu)$ and $|\mathcal{U}(u_2) \cap C(wu)| \leq \Delta - 2$; similarly, we also have that $|\mathcal{U}(u_3) \cap C(wu)| \leq \Delta - 2$ and $|\mathcal{U}(u_4) \cap C(wu)| \leq \Delta - 2$.

Suppose that the color 1 appears at most twice in $\mathbb{S}$; by symmetry, assume that $1 \notin \mathcal{U}(u_3) \cup \mathcal{U}(u_4)$. Thus there exists a $(2, 1, u, u_4)$-critical path; otherwise, reassigning 1 to uu_4 will take us back to Subcase 2.2.2. But reassigning 1 to uu_3 will take us back to Subcase 2.2.2 again. Hence, the color 1 appears at least three times in $\mathbb{S}$.

Furthermore, $\mathcal{A}(u_2) \cup \mathcal{A}(u_3) \cup \mathcal{A}(u_4) \nsubseteq \mathcal{U}(w_1)$; otherwise, we have $|\mathcal{U}(w_1)| \geq 2 + |\mathcal{A}(u_2)| + |\mathcal{A}(u_3)| + |\mathcal{A}(u_4)| > \deg_G(w_1)$, which is a contradiction. Without loss of generality, assume that $\beta_1 \notin \mathcal{U}(w_1)$. Clearly, there exists a $(3, \beta_1, u, w)$- or $(4, \beta_1, u, w)$-critical path. By symmetry, assume that there exists a $(3, \beta_1, u, w)$-critical path. There exists a $(4, \beta_1, u, u_2)$-critical path; otherwise, reassigning β_1 to uu_2 will take us back to Subcase 2.1. It follows that $4 \in \mathcal{U}(u_2)$.

If $2 \notin \mathbb{S}$, then reassigning 2 to uu_4 and β_1 to uu_2 will take us back to Subcase 2.1. So we have $2 \in \mathbb{S}$; similarly, we can obtain that $\mathcal{U}(w) \setminus \{1, 3, 4\} \subseteq \mathbb{S}$.

If $3 \notin \mathbb{S}$, then $4 \in \mathcal{U}(w_1) \cup \mathcal{U}(u_3)$; otherwise, reassigning $3, 4$ and β_1 to uu_4, uu_3 and uu_2 respectively, and then we go back to Subcase 2.1. Anyway, we have that $\mathrm{mul}_{\mathbb{S}}(3) + \mathrm{mul}_{\mathbb{S}}(4) \geq 2$.

There exists a $(2, \kappa - \Delta, u, w)$- or $(3, \kappa - \Delta, u, w)$- or $(4, \kappa - \Delta, u, w)$-critical path; otherwise, reassigning β_1 to uw_1 and $\kappa - \Delta$ to uw results in an acyclic edge coloring of G. If $\kappa - \Delta \notin \mathcal{U}(u_3) \cup \mathcal{U}(u_4)$, then reassigning $\kappa - \Delta$ to uu_3 and β_1 to uw_1 will take us back to Case 2.1. This implies that $\kappa - \Delta \in \mathcal{U}(u_3) \cup \mathcal{U}(u_4)$; similarly, we can prove that $\kappa - \Delta \in \mathcal{U}(u_2) \cup \mathcal{U}(u_4)$ and $\kappa - \Delta \in \mathcal{U}(u_2) \cup \mathcal{U}(u_3)$. Hence, the color $\kappa - \Delta$ appears at least twice in $\mathbb{S}$. Therefore, we have

$$\sum_{\theta \in \mathcal{U}(w) \cup \mathcal{U}(u)} \mathrm{mul}_{\mathbb{S}}(\theta) + |X| \geq 3|\{1\}| + 2|\{\kappa - \Delta\}| + \sum_{\theta \in \mathcal{U}(w) \setminus \{1\}} \mathrm{mul}_{\mathbb{S}}(\theta) \geq \deg_G(w) + 3.$$

Subcase 2.3. $|\mathcal{U}(w) \cap \mathcal{U}(u)| = 4$.

In other words, $\mathcal{U}(u) \subseteq \mathcal{U}(w)$. It follows that $|C(wu)| = \kappa - \deg_G(w) + 1 \geq \Delta + 1$ and $|\mathcal{A}(u_i)| \geq 2$ for $i = 2, 3, 4$. By Claim 2, we have that $\mathcal{A}(u_2), \mathcal{A}(u_3)$ and $\mathcal{A}(u_4)$ are pairwise disjoint and $\mathcal{U}(w_1) \supseteq \mathcal{A}(u_2) \cup \mathcal{A}(u_3) \cup \mathcal{A}(u_4)$, which implies that $|\mathcal{U}(w_1)| \geq 2 + |\mathcal{A}(u_2)| + |\mathcal{A}(u_3)| + |\mathcal{A}(u_4)| > \deg_G(w_1)$, a contradiction. □

4 The main result

Now, we are ready to prove the main result, Theorem 1.1.



(a) $(3, 9^+, 9^+)$-face



(b) $(4, 4, 10^+)$-face



(c) $(4, 5, 11)$-face



(d) $(4, 5, \neq 11)$-face



(e) $(4, 6, 10^+)$-face



(f) $(4, 7–8, 7–9)$-face



(g) $(4, 7–8, 10^+)$-face



(h) $(4, 9^+, 9^+)$-face



(i) $(4, 9^+, 9^+)$-face



(j) $(4, 9^+, 9^+)$-face



(k) $(5, 5, 5–7)$-face



(l) $(5, 5, 8^+)$-face



(m) $(5, 6, 6)$-face



(n) $(5, 6, 7)$-face



(o) $(5, 6, 8^+)$-face



(p) $(5, 7, 7)$-face



(q) $(5, 7, 8^+)$-face



(r) $(5, 8^+, 8^+)$-face



(s) $(6^+, 6^+, 6^+)$-face

Fig. 1: Discharging rules

Proof of Theorem 1.1. Suppose that G is a counterexample with $|V| + |E|$ is minimum, and fix $\kappa = \Delta(G) + 6$. Since the hypothesis is minor-closed, it follows that G is a κ-minimal graph. Let G^* be obtained from G by removing all the 2-vertices. By Lemma 1 and Lemma 3, the minimum degree of G^* is at least three. Take a component H of G^* and embed it in the plane. In the following, we will do arguments on the graph H to obtain a contradiction.

By Lemma 3 (A), we have the following claims.

Claim 1. If $\deg_H(v) < \deg_G(v)$, then $\deg_H(v) \geq 8 + m$, where m is the number of adjacent 7^--vertices in H.

Claim 2. If $\deg_H(v) \leq 7$, then $\deg_G(v) = \deg_H(v)$.

From the Euler's formula, we have the following equality:

$$\sum_{v \in V(H)} (2\deg_H(v) - 6) + \sum_{f \in F(H)} (\deg_H(f) - 6) = -12 \tag{8}$$

Assign the initial charge of every vertex v to be $2\deg_H(v) - 6$ and the initial charge of every face f to be $\deg_H(f) - 6$. Clearly, the sum of the initial charge of vertices and faces is -12. We design appropriate discharging rules and redistribute charge among the vertices and faces, such that the final charge of every vertex and every face is nonnegative, which derive a contradiction.

Discharging Rules:

(R1) If w is a 4-vertex adjacent to a 5^--vertex u, then w sends $\frac{4}{5}$ to each face incident with wu, and sends $\frac{1}{5}$ to each other face.

(R2) If w is a 4-vertex adjacent to a 6-vertex u, then w sends $\frac{2}{3}$ to each face incident with wu, and sends $\frac{1}{3}$ to each other face.

(R3) If w is a 4-vertex which is not adjacent to 6^--vertices, then w sends $\frac{1}{2}$ to each incident face.

(R4) All the rules regarding 3-faces are in the Fig (a)–(s).

(R5) Every 9^+-vertex sends 1 to each incident 4^+-face.

(R6) Every vertex with degree $5, 6, 7$ or 8 sends $\frac{1}{2}$ to each incident 4^+-face.

Computing the final charge of faces.

Let $f = w_1w_2w_3$ be a 3-face with $\deg_H(w_1) \leq \deg_H(w_2) \leq \deg_H(w_3)$.

If w_1 is a 3-vertex, then Lemma 6 implies that both w_2 and w_3 are 9^+-vertices in G, and they also are 9^+-vertices in H by Claim 1, thus f is a $(3, 9^+, 9^+)$-face in H and the final charge is $-3 + 2 \times \frac{3}{2} = 0$.

If w_1w_2 is a $(4,4)$-edge, then Lemma 9 implies that w_3 is a 12^+-vertex in G, and it is a 10^+-vertex in H by Claim 1, thus f is a $(4, 4, 10^+)$-face and the final charge is $-3 + 2 \times \frac{4}{5} + \frac{7}{5} = 0$.

If w_1w_2 is a $(4,5)$-edge, then Lemma 9 implies that w_3 is a 11^+-vertex in G, and it is a 10^+-vertex in H by Claim 1, thus the final charge of f is $-3 + \frac{4}{5} + \frac{17}{20} + \frac{27}{20} = 0$ if $\deg_H(w_3) = 11$, or $-3 + 2 \times \frac{4}{5} + \frac{7}{5} = 0$ if w_3 is a 10- or 12^+-vertex in H.

If w_1w_2 is a $(4,6)$-edge, then Lemma 9 implies that w_3 is a 10^+-vertex in G, and it is a 10^+-vertex in H by Claim 1, and then the final charge is $-3 + \frac{2}{3} + 1 + \frac{4}{3} = 0$.

If $\deg_H(w_1) = 4$, $\deg_H(w_2) \in \{7, 8\}$ and $\deg_H(w_3) \in \{7, 8, 9\}$, then the final charge of f is $-3 + \frac{1}{2} + 2 \times \frac{5}{4} = 0$.

If $\deg_H(w_1) = 4$, $\deg_H(w_2) \in \{7, 8\}$ and $\deg_H(w_3) \geq 10$, then the final charge of f is $-3 + \frac{1}{2} + \frac{7}{6} + \frac{4}{3} = 0$.

Suppose that f is a $(4, 9^+, 9^+)$-face. If w_1 is adjacent to a 5^--vertex u, then w_1 sends $\frac{1}{5}$ to f, and then the final charge of f is $-3 + \frac{1}{5} + 2 \times \frac{7}{5} = 0$; if w_1 is adjacent to a 6-vertex u, then w_1 sends $\frac{1}{3}$ to f, and then the final charge of f is $-3 + \frac{1}{3} + 2 \times \frac{4}{3} = 0$; if w_1 is not adjacent to 6^--vertices, then w_1 sends $\frac{1}{2}$ to f, and then the final charge of f is $-3 + \frac{1}{2} + 2 \times \frac{5}{4} = 0$.

If $\deg_H(w_1) = \deg_H(w_2) = 5$ and $\deg_H(w_3) \in \{5, 6, 7\}$, then the final charge of f is $-3 + 3 \times 1 = 0$.

If f is a $(5, 5, 8^+)$-face, then the final charge is $-3 + 2 \times \frac{7}{8} + \frac{5}{4} = 0$.

If f is a $(5, 6, 6)$-face, then the final charge is $-3 + 3 \times 1 = 0$.

If f is a $(5, 6, 7)$-face, then the final charge is $-3 + \frac{5}{6} + 1 + \frac{7}{6} = 0$.

If f is a $(5,6,8^+)$-face, then the final charge is $-3+\frac{3}{4}+1+\frac{5}{4}=0$.
If f is a $(5,7,7)$-face, then the final charge is $-3+\frac{2}{3}+2\times\frac{7}{6}=0$.
If f is a $(5,7,8^+)$-face, then the final charge is $-3+\frac{17}{28}+\frac{8}{7}+\frac{5}{4}=0$.
If f is a $(5,8^+,8^+)$-face, then the final charge is $-3+\frac{1}{2}+2\times\frac{5}{4}=0$.
If f is a $(6^+,6^+,6^+)$-face, then the final charge is $-3+3\times 1=0$.

Next, we compute the final charge of 4-faces. Let $w_1w_2w_3w_4$ be a 4-face with w_2 having the minimum degree on the boundary. If $\deg_H(w_2)\geq 5$, then the final charge of f is at least $-2+4\times\frac{1}{2}=0$. If $\deg_H(w_1),\deg_H(w_3)\geq 9$, then the final charge is at least $-2+2\times 1=0$. So we may assume that $\deg_H(w_2)\in\{3,4\}$ and $\deg_H(w_1)\leq 8$. By Lemma 6 and Claim 1, we have that $\deg_H(w_2)=4$ and $\deg_G(w_1)=\deg_H(w_1)\leq 8$. By Lemma 9 and discharging rules, the face f receives at least $\frac{1}{2}$ from each incident vertex, so the final charge of f is at least $-2+4\times\frac{1}{2}=0$.

Suppose that f is a 5-face. If f is incident with a 9^+-vertex, then the final charge is at least $-1+1=0$. So we may assume that f is incident with five 8^--vertices. It is obvious that f is incident with at least two 5^+-vertices, and then the final charge is at least $-1+2\times\frac{1}{2}=0$.

If f is a 6^+-face, then the final charge is at least $\deg_H(f)-6\geq 0$.

Computing the final charge of vertices.

Let v be a 3-vertex. Clearly, the final charge is zero.

Let v be a 4-vertex. If v is adjacent to a 5^--vertex, then Lemma 9 and Claim 1 implies that v is adjacent to three 9^+-vertices, and then the final charge is $2-2\times\frac{4}{5}-2\times\frac{1}{5}=0$. If v is adjacent to a 6-vertex, then Lemma 9 and Claim 1 implies that v is adjacent to three 9^+-vertices, and then the final charge is $2-2\times\frac{2}{3}-2\times\frac{1}{3}=0$. If v is not adjacent to 6^--vertices, then the final charge is $2-4\times\frac{1}{2}=0$.

Let v be a 5-vertex with neighbors $v_1,v_2,\ldots,v_5$ in anticlockwise order. If v sends at most $\frac{4}{5}$ to each incident face, then the final charge is at least $4-5\times\frac{4}{5}=0$. So we may assume that v sends more than $\frac{4}{5}$ to some face f.

If f is a $(5,5,5)$-face, then Lemma 10 and Claim 1 implies that the other three vertices adjacent to v are 9^+-vertices, and then the final charge of v is at least $4-1-2\times\frac{7}{8}-2\times\frac{1}{2}>0$.

If f is a $(5,5,6)$-face, then Lemma 10 and Claim 1 implies that the other three vertices adjacent to v are 8^+-vertices, and then the final charge of v is at least $4-1-\frac{7}{8}-\frac{3}{4}-2\times\frac{1}{2}>0$.

If f is a $(5,5,7)$-face, then Lemma 10 and Claim 1 implies that the other three vertices adjacent to v are 7^+-vertices, and then the final charge of v is at least $4-2\times 1-3\times\frac{2}{3}=0$.

If f is a $(5,6,6)$-face, then Lemma 10 and Claim 1 implies that the other three vertices adjacent to v are 7^+-vertices, and then the final charge of v is at least $4-1-2\times\frac{5}{6}-2\times\frac{2}{3}=0$.

If v sends at most $\frac{1}{2}$ to an incident face, then the final charge of v is at least $4-4\times\frac{7}{8}-\frac{1}{2}=0$. So we may assume that the 5-vertex v sends more than $\frac{1}{2}$ to each incident face, thus v is incident with five 3-faces.

Suppose that $f=vv_1v_2$ is a 3-face with $\deg_H(v_1)=5$ and $\deg_H(v_2)\geq 8$. By the excluded cases in the above, the vertex v_5 is an 8^+-vertex. Since v sends more than $\frac{1}{2}$ to the 3-face vv_2v_3, the vertex v_3 is a 7^--vertex. Similarly, the vertex v_4 is also a 7^--vertex. Now, the 3-face vv_3v_4 is a $(5,7^-,7^-)$-face. By the excluded cases, we only have to consider the edge v_3v_4 is a $(6,7)$- or $(7,6)$- or $(7,7)$-edge. If v_3v_4 is a $(7,7)$-edge, then the final charge of v is at least $4-2\times\frac{7}{8}-2\times\frac{17}{28}-\frac{2}{3}>0$. If v_3v_4 is $(6,7)$- or $(7,6)$-edge, then the final charge of v is at least $4-2\times\frac{7}{8}-\frac{3}{4}-\frac{5}{6}-\frac{17}{28}>0$.

Suppose that $f=vv_1v_2$ is a $(5,6,7)$-face with $\deg_H(v_1)=6$ and $\deg_H(v_2)=7$. By the excluded cases, the vertex v_3 is a 6^+-vertex and the vertex v_5 is a 7^+-vertex. By Lemma 6 and Claim 1, the vertex v_4 is a 4^+-vertex. If $\deg_H(v_4)=4$, then Lemma 9 and Claim 1 implies that both v_3 and v_5 are 11^+-vertices, thus the final charge of v is at least $4-\frac{5}{6}-\frac{3}{4}-\frac{17}{28}-2\times\frac{17}{20}>0$. By the excluded cases, the vertex v_4 cannot be a 5-vertex. If $\deg_H(v_4)=6$, then $\deg_H(v_3)\geq 7$, and then the final charge of v is at least $4-\frac{2}{3}-4\times\frac{5}{6}=0$. If $\deg_H(v_4)\geq 7$, then the final charge is at least $4-\frac{2}{3}-4\times\frac{5}{6}=0$.

Suppose that $f=vv_1v_2$ is a $(5,4,11)$-face. By Lemma 9 and Claim 1, the vertex v_5 is a 10^+-vertex. If one of v_3 and v_4 is a 8^+-vertex, then v sends $\frac{1}{2}$ to an incident 3-face, a contradiction. So we may assume that $\deg_H(v_3),\deg_H(v_4)\leq 7$. By the excluded cases, the edge v_3v_4 is a $(7,7)$-edge, and then the final charge of v is at least $4-2\times\frac{17}{28}-\frac{2}{3}-2\times\frac{17}{20}>0$.

Let v be a 6-vertex. The final charge is at least $6-6\times 1=0$.

Let v be a 7-vertex. If v sends at most $\frac{1}{2}$ to an incident face, then the final charge is at least $8-6\times\frac{5}{4}-\frac{1}{2}=0$. So we may assume that v sends more than $\frac{1}{2}$ to each incident face, thus v is incident with seven 3-faces. By Lemma 9 (b) and Claim 1, the vertex v is not incident with $(4,7,9^-)$-faces. Now, the vertex v sends at most $\frac{7}{6}$ to each incident face. If v is incident with a $(5^-,5^-,7)$- or $(6^+,6^+,7)$-face, then the final charge is at least $8-6\times\frac{7}{6}-1=0$. So every face incident with v is a $(5^-,6^+,7)$-face, but the vertex v is a 7-vertex and the number 7 is odd, a contradiction.

Let v be an 8-vertex. Every 8-vertex sends at most $\frac{5}{4}$ to each incident face, thus the final charge is at least $10-8\times\frac{5}{4}=0$.

Let v be a 9-vertex. If $\deg_G(v)>9$, then Claim 1 implies that v is adjacent to at most one 7^--vertex in H, and then the final charge of v is at least $12-7\times1-2\times\frac{3}{2}>0$. So we may assume that $\deg_G(v)=\deg_H(v)=9$.

Suppose that $(3,9)$-edge uv is incident with two 3-faces. By Lemma 7, the vertex v is adjacent to eight 8^+-vertices, and then the final charge is at least $12-7\times1-2\times\frac{3}{2}>0$. So every $(3,9)$-edge uv is incident with at most one 3-face.

Let τ be the number of incident 4^+-faces. If $\tau\geq4$, then the final charge is at least $12-5\times\frac{3}{2}-4\times1>0$. Since $\deg_G(v)=\deg_H(v)=9$, Lemma 9 implies that v is not incident with face (h) or (i). If $\tau\leq3$, then the final charge is at least $12-\tau-2\tau\times\frac{3}{2}-(9-3\tau)\times\frac{5}{4}\geq0$.

Let v be a 10-vertex. If $\deg_G(v)>10$, then Claim 1 implies that v is adjacent to at most two 7^--vertices, and then the final charge is at least $14-4\times\frac{3}{2}-6\times1>0$. So we may assume that $\deg_G(v)=\deg_H(v)=10$. Hence, the vertex v is not incident with face (b), (d) or (h), and thus v sends $\frac{3}{2}$, $\frac{4}{3}$, $\frac{5}{4}$ or 1 to each incident face.

If v is incident with at least two 4^+-faces, then the final charge is at least $14-8\times\frac{3}{2}-2\times1=0$. Hence, the vertex v is incident with at most one 4^+-face. Lemma 9 implies that v is adjacent to at most five 4^--vertices. Let s be the number of incident $(10,3,9^+)$-faces, and let s^* be the number of incident $(10,4,6^+)$-faces.

If $s\leq4$, then the final charge is at least $14-s\times\frac{3}{2}-(10-s)\times\frac{4}{3}=\frac{2}{3}-\frac{s}{6}\geq0$. So we may assume that $s\geq5$, and then the number of adjacent 3-vertices is at least three.

(1) $s\in\{5,6\}$.

If $s^*=0$, then the final charge is at least $14-6\times\frac{3}{2}-4\times\frac{5}{4}=0$. If v is incident with exactly one 4^+-face, then the final charge is at least $14-6\times\frac{3}{2}-1-3\times\frac{4}{3}=0$. So we may assume that $s^*\geq1$ and v is not incident with any 4^+-face. Clearly, the vertex v is incident with exactly six $(10,3,9^+)$-faces and $s=6$. It is obvious that v is adjacent to at least one 4-vertex. Lemma 8 implies that the vertex v is adjacent to exactly three 3-vertices, one 4-vertex and six 6^+-vertices. Hence, it is incident with exactly two $(10,6^+,6^+)$-faces, and then the final charge is at least $14-6\times\frac{3}{2}-2\times\frac{4}{3}-2\times1>0$.

(2) $s\geq7$.

Clearly, the vertex v is adjacent to at least four 3-vertices. Lemma 8 implies that the vertex v is adjacent to exactly four 3-vertices and six 6^+-vertices. Hence, the vertex v is incident with two $(10,6^+,6^+)$-faces, or one $(10,6^+,6^+)$-face and one 4^+-face, thus the final charge is at least $14-8\times\frac{3}{2}-2\times1=0$.

Let v be an 11-vertex. If $\deg_G(v)>11$, then v is adjacent to at most three 7^--vertices in H, and then the final charge is at least $16-6\times\frac{3}{2}-5\times1>0$. So we may assume that $\deg_G(v)=\deg_H(v)=11$.

If v sends at most 1 to an incident face, then the final charge is at least $16-10\times\frac{3}{2}-1=0$. So we may assume that v is not incident with 4^+-faces and is not incident with $(11,6^+,6^+)$-faces. Since the degree of v is odd, the vertex v cannot be incident with eleven $(11,5^-,6^+)$-faces. So v is incident with a $(11,5^-,5^-)$-face f. Lemma 6 and Lemma 9 implies that the face f is a $(4,5,11)$-face or $(5,5,11)$-face. Hence, the vertex v is adjacent to at most four 3-vertices. If v is adjacent to at most three 3-vertices, then the final charge is at least $16-6\times\frac{3}{2}-5\times\frac{7}{5}=0$. Hence, the vertex v is adjacent to exactly four 3-vertices, see Fig. 2. If f is a $(5,5,11)$-face, then the final charge of v is $16-8\times\frac{3}{2}-3\times\frac{5}{4}>0$. If f is a $(4,5,11)$-face, then the final charge is $16-8\times\frac{3}{2}-\frac{5}{4}-\frac{27}{20}-\frac{7}{5}=0$.

Let v be a 12^+-vertex. The final charge is at least $2\deg_H(v)-6-\deg_H(v)\times\frac{3}{2}=\frac{1}{2}\deg_H(v)-6\geq0$. □



Fig. 2: The vertex x is a 4- or 5-vertex.

Acknowledgments. This project was supported by the National Natural Science Foundation of China (11101125) and partially supported by the Fundamental Research Funds for Universities in Henan. The authors would like to thank the referees for their valuable comments.

References

[1] N. Alon, C. McDiarmid and B. Reed, Acyclic coloring of graphs, Random Structures Algorithms 2 (3) (1991) 277–288.

[2] N. Alon, B. Sudakov and A. Zaks, Acyclic edge colorings of graphs, J. Graph Theory 37 (3) (2001) 157–167.

[3] M. Basavaraju, L. S. Chandran, N. Cohen, F. Havet and T. Müller, Acyclic edge-coloring of planar graphs, SIAM J. Discrete Math. 25 (2) (2011) 463–478.

[4] L. Esperet and A. Parreau, Acyclic edge-coloring using entropy compression, European J. Combin. 34 (6) (2013) 1019–1027.

[5] I. Fiamčík, The acyclic chromatic class of a graph, Math. Slovaca 28 (2) (1978) 139–145.

[6] A. Fiedorowicz, M. Hałuszczak and N. Narayanan, About acyclic edge colourings of planar graphs, Inform. Process. Lett. 108 (6) (2008) 412–417.

[7] I. Giotis, L. Kirousis, K. I. Psaromiligkos and D. M. Thilikos, On the algorithmic Lovász local lemma and acyclic edge coloring, eprint arXiv:1407.5374.

[8] Y. Guan, J. Hou and Y. Yang, An improved bound on acyclic chromatic index of planar graphs, Discrete Math. 313 (10) (2013) 1098–1103.

[9] J. Hou, N. Roussel and J. Wu, Acyclic chromatic index of planar graphs with triangles, Inform. Process. Lett. 111 (17) (2011) 836–840.

[10] J. Hou, J. Wu, G. Liu and B. Liu, Acyclic edge colorings of planar graphs and series-parallel graphs, Sci. China Ser. A 52 (3) (2009) 605–616.

[11] M. Molloy and B. Reed, Further algorithmic aspects of the local lemma, in: Proceedings of the Thirtieth Annual ACM Symposium on the Theory of Computing, ACM, New York, 1998, pp. 524–529.

[12] S. Ndreca, A. Procacci and B. Scoppola, Improved bounds on coloring of graphs, European J. Combin. 33 (4) (2012) 592–609.

[13] T. Wang and Y. Zhang, Acyclic edge coloring of graphs, Discrete Appl. Math. 167 (2014) 290–303.

[14] W. Wang, Q. Shu and Y. Wang, A new upper bound on the acyclic chromatic indices of planar graphs, European J. Combin. 34 (2) (2013) 338–354.